

L Number	Hits	Search Text	DB	Time stamp
1	10252	("514/183,247,272,277,315,318,333,336,340,364,383").CCLS	USPAT	2003/10/13 11:21
2	3737	("546/184,193,210,223,236,269,1").CCLS	USPAT	2003/10/13 11:21
3	779	("548/125,131").CCLS	USPAT	2003/10/13 11:22
4	51	((("514/183,247,272,277,315,318,333,336,340,364,383").CCLS) and (("546/184,193,210,223,236,269,1").CCLS) and ((("548/125,131").CCLS)	USPAT	2003/10/13 11:22

L Number	Hits	Search Text	DB	Time stamp
1	10252	("514/183, 247, 272, 277, 315, 318, 333, 336, 340, 364, 383").CCLS	USPAT	2003/10/13 11:21
2	3737	("546/184, 193, 210, 223, 236, 269, 1").CCLS	USPAT	2003/10/13 11:21
3	779	("548/125, 131").CCLS	USPAT	2003/10/13 11:22
4	51	(("514/183, 247, 272, 277, 315, 318, 333, 336, 340, 364, 383").CCLS) and ((("546/184, 193, 210, 223, 236, 269, 1").CCLS) and ((("548/125, 131").CCLS)	USPAT	2003/10/13 11:22

L Number	Hits	Search Text	DB	Time stamp
1	10252	("514/183, 247, 272, 277, 315, 318, 333, 336, 340, 364, 383") .CCLS	USPAT	2003/10/13 11:21
2	3737	("546/184, 193, 210, 223, 236, 269.1") .CCLS	USPAT	2003/10/13 11:21
3	779	("548/125, 131") .CCLS	USPAT	2003/10/13 11:22
4	51	((("514/183, 247, 272, 277, 315, 318, 333, 336, 340, 364, 383") .CCLS) and (("546/184, 193, 210, 223, 236, 269.1") .CCLS) and ((("548/125, 131") .CCLS)	USPAT	2003/10/13 11:25
5	7	((("514/183, 247, 272, 277, 315, 318, 333, 336, 340, 364, 383") .CCLS) and (("546/184, 193, 210, 223, 236, 269.1") .CCLS) and ((("548/125, 131") .CCLS)) and piperidine and 1,2,4-oxadiazole	USPAT	2003/10/13 11:25

L Number	Hits	Search Text	DB	Time stamp
1	10252	("514/183,247,272,277,315,318,333,336,340,364,383").CCLS	USPAT	2003/10/13 11:21
2	3737	("546/184,193,210,223,236,269,1").CCLS	USPAT	2003/10/13 11:21
3	779	("548/125,131").CCLS	USPAT	2003/10/13 11:22
4	51	(("514/183,247,272,277,315,318,333,336,340,364,383") .CCLS) and (("546/184,193,210,223,236,269,1") .CCLS) and (("548/125,131") .CCLS)	USPAT	2003/10/13 11:25
5	7	((("514/183,247,272,277,315,318,333,336,340,364,383") .CCLS) and (("546/184,193,210,223,236,269,1") .CCLS) and (("548/125,131") .CCLS)) and piperidine and 1,2,4-oxadiazole	USPAT	2003/10/13 11:25

Welcome to STN International! Enter x:x

LOGINID:ssspta1611sxp

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * * * * * Welcome to STN International * * * * * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 SEP 09 CA/CAplus records now contain indexing from 1907 to the present
NEWS 4 AUG 05 New pricing for EUROPATFULL and PCTFULL effective August 1, 2003
NEWS 5 AUG 13 Field Availability (/FA) field enhanced in BEILSTEIN
NEWS 6 AUG 18 Data available for download as a PDF in RDISCLOSURE
NEWS 7 AUG 18 Simultaneous left and right truncation added to PASCAL
NEWS 8 AUG 18 FROSTI and KOSMET enhanced with Simultaneous Left and Right Truncation
NEWS 9 AUG 18 Simultaneous left and right truncation added to ANABSTR
NEWS 10 SEP 22 DIPPR file reloaded
NEWS 11 SEP 25 INPADOC: Legal Status data to be reloaded
NEWS 12 SEP 29 DISSABS now available on STN
NEWS 13 OCT 10 PCTFULL: Two new display fields added

NEWS EXPRESS OCTOBER 01 CURRENT WINDOWS VERSION IS V6.01a, CURRENT MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP), AND CURRENT DISCOVER FILE IS DATED 23 SEPTEMBER 2003

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * * * * * STN Columbus * * * * * * * * * * * * *

FILE 'HOME' ENTERED AT 10:39:40 ON 13 OCT 2003

=> file reg	COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST		0.21	0.21

FILE 'REGISTRY' ENTERED AT 10:39:56 ON 13 OCT 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 12 OCT 2003 HIGHEST RN 603065-76-5
DICTIONARY FILE UPDATES: 12 OCT 2003 HIGHEST RN 603065-76-5

TSCA INFORMATION NOW CURRENT THROUGH JULY 14, 2003

Please note that search-term pricing does apply when
conducting SmartSELECT searches.


Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>
Uploading 10069215.1

L1 STRUCTURE UPLOADED

=> d 11
L1 HAS NO ANSWERS
L1 STR



G1 O,N,C,S,CH

Structure attributes must be viewed using STN Express query preparation.

=> s 11 sss full
 FULL SEARCH INITIATED 10:40:24 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 377 TO ITERATE

100.0% PROCESSED 377 ITERATIONS 1 ANSWERS
 SEARCH TIME: 00.00.01

L2 1 SEA SSS FUL L1

=> file marpat
 COST IN U.S. DOLLARS SINCE FILE TOTAL
 ENTRY SESSION
 FULL ESTIMATED COST 148.15 148.36

FILE 'MARPAT' ENTERED AT 10:40:31 ON 13 OCT 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 American Chemical Society (ACS)

FILE CONTENT: 1988-PRESENT (VOL 104 ISS 15-VOL 139 ISS15) (20031010ED)

MOST RECENT CITATIONS FOR PATENTS FROM FIVE MAJOR ISSUING AGENCIES
 (COVERAGE TO THESE DATES IS NOT COMPLETE) :

US 6617454 09 SEP 2003
 DE 20300703 18 SEP 2003
 EP 1342759 10 SEP 2003
 JP 2003261500 16 SEP 2003
 WO 2003076638 18 SEP 2003

Structure search limits have been raised. See HELP SLIMIT for the new,
 higher limits.

=> s 11 sss full
 STRUCTURE TOO LARGE - SEARCH ENDED
 A structure in your query is too large. You may delete
 attributes or atoms to reduce the size of the structure
 and try again.

=> file caold
 COST IN U.S. DOLLARS SINCE FILE TOTAL
 ENTRY SESSION
 FULL ESTIMATED COST 0.40 148.76

FILE 'CAOLD' ENTERED AT 10:41:01 ON 13 OCT 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1907-1966
 FILE LAST UPDATED: 01 May 1997 (19970501/UP)

This file contains CAS Registry Numbers for easy and accurate
 substance identification. Title keywords, authors, patent
 assignees, and patent information, e.g., patent numbers, are
 now searchable from 1907-1966. TIFF images of CA abstracts
 printed between 1907-1966 are available in the PAGE

display formats.

This file supports REG1stRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

=> s ll sss full

REG1stRY INITIATED

Substance data SEARCH and crossover from CAS REGISTRY in progress...
Use DISPLAY HITSTR (or FHITSTR) to directly view retrieved structures.

FULL SEARCH INITIATED 10:41:08 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 377 TO ITERATE

100.0% PROCESSED 377 ITERATIONS
SEARCH TIME: 00.00.01

1 ANSWERS

L3 1 SEA SSS FUL L1

L4 0 L3

=> file caplus
COST IN U.S. DOLLARS SINCE FILE TOTAL
FULL ESTIMATED COST ENTRY SESSION
0.40 297.71

FILE 'CAPLUS' ENTERED AT 10:41:26 ON 13 OCT 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 Oct 2003 VOL 139 ISS 16
FILE LAST UPDATED: 12 Oct 2003 (20031012/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d his

(FILE 'HOME' ENTERED AT 10:39:40 ON 13 OCT 2003)

```
FILE 'REGISTRY' ENTERED AT 10:39:56 ON 13 OCT 2003
L1      STRUCTURE uploaded
L2      1 S L1 SSS FULL

FILE 'MARPAT' ENTERED AT 10:40:31 ON 13 OCT 2003

FILE 'CAOLD' ENTERED AT 10:41:01 ON 13 OCT 2003
S L1

FILE 'REGISTRY' ENTERED AT 10:41:07 ON 13 OCT 2003
L3      1 S L1 SSS FULL

FILE 'CAOLD' ENTERED AT 10:41:09 ON 13 OCT 2003
L4      0 S L3 SSS FULL

FILE 'CAPLUS' ENTERED AT 10:41:26 ON 13 OCT 2003

=> s 12
L5      2 L2

=> d 12 fbib hitstr abs total
YOU HAVE REQUESTED DATA FROM FILE 'REGISTRY' - CONTINUE? (Y)/N:y
```

'FBIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'
'HITSTR' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'
'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = 'CAS Registry Number')

REG	- RN
SAM	- Index Name, MF, and structure - no RN
FIDE	- All substance data, except sequence data
IDE	- FIDE, but only 50 names
SQIDE	- IDE, plus sequence data
SQIDE3	- Same as SQIDE, but 3-letter amino acid codes are used
SQD	- Protein sequence data, includes RN
SQD3	- Same as SQD, but 3-letter amino acid codes are used
SQN	- Protein sequence name information, includes RN
CALC	- Table of calculated properties
EPROP	- Table of experimental properties
PROP	- EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS	-- Abstract
APPS	-- Application and Priority Information
BIB	-- CA Accession Number, plus Bibliographic Data
CAN	-- CA Accession Number
CBIB	-- CA Accession Number, plus Bibliographic Data (compressed)

IND -- Index Data
 IPC -- International Patent Classification
 PATS -- PI, SO
 STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels
 IBIB -- BIB, indented, with text labels
 ISTD -- STD format, indented
 OBIB ----- AN, plus Bibliographic Data (original)
 OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
 SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.


The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.
 HELP FORMATS -- To see detailed descriptions of the predefined formats.
 ENTER DISPLAY FORMAT (IDE):IDE

L2 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS on STN
 RN 328083-09-6 REGISTRY
 CN 1,2,4-Oxadiazole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-(2-pyridinyl)- (9CI) (CA INDEX NAME)
 OTHER NAMES:
 CN N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[3-(2-pyridinyl)-1,2,4-oxadiazol-5-yl]acetamide
 FS 3D CONCORD
 MF C21 H21 Cl2 N5 O2
 SR CA
 LC STN Files: CA, CAPLUS



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> file caplus	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.42	300.63

FILE 'CAPLUS' ENTERED AT 10:42:33 ON 13 OCT 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 Oct 2003 VOL 139 ISS 16
 FILE LAST UPDATED: 12 Oct 2003 (20031012/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 11
REGISTRY INITIATED
 Substance data SEARCH and crossover from CAS REGISTRY in progress...
 Use DISPLAY HITSTR (or FHITSTR) to directly view retrieved structures.

SAMPLE SEARCH INITIATED 10:42:46 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 16 TO ITERATE

100.0% PROCESSED 16 ITERATIONS 0 ANSWERS
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 80 TO 560
 PROJECTED ANSWERS: 0 TO 0

L6 0 SEA SSS SAM L1

L7 0 L6

=> d his

(FILE 'HOME' ENTERED AT 10:39:40 ON 13 OCT 2003)

FILE 'REGISTRY' ENTERED AT 10:39:56 ON 13 OCT 2003
L1 STRUCTURE uploaded
L2 1 S L1 SSS FULL

FILE 'MARPAT' ENTERED AT 10:40:31 ON 13 OCT 2003

FILE 'CAOLD' ENTERED AT 10:41:01 ON 13 OCT 2003
S L1

FILE 'REGISTRY' ENTERED AT 10:41:07 ON 13 OCT 2003
L3 1 S L1 SSS FULL

FILE 'CAOLD' ENTERED AT 10:41:09 ON 13 OCT 2003
L4 0 S L3 SSS FULL

FILE 'CPLUS' ENTERED AT 10:41:26 ON 13 OCT 2003
L5 2 S L2

FILE 'REGISTRY' ENTERED AT 10:42:19 ON 13 OCT 2003

FILE 'CPLUS' ENTERED AT 10:42:26 ON 13 OCT 2003

FILE 'CPLUS' ENTERED AT 10:42:33 ON 13 OCT 2003
S L1

FILE 'REGISTRY' ENTERED AT 10:42:45 ON 13 OCT 2003
L6 0 S L1

FILE 'CPLUS' ENTERED AT 10:42:47 ON 13 OCT 2003
L7 0 S L6

=> file caplus	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.42	301.87

FILE 'CPLUS' ENTERED AT 10:43:40 ON 13 OCT 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 Oct 2003 VOL 139 ISS 16
FILE LAST UPDATED: 12 Oct 2003 (20031012/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.


=> S 15
 L8 2 L2

=> D 18 fbib hitstr abs total



L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:44146 CAPLUS
 DN 138:73178
 TI Preparation and pharmaceutical combinations of
 [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3 antagonists
 for treatment of asthma, allergic disease, or inflammation
 IN Bahl, Ash; Perry, Matthew; Springthorpe, Brian
 PA AstraZeneca AB, Swed.
 SO Brit. UK Pat. Appl., 91 pp.
 CODEN: BAXXDU
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 2373186	A1	20020918	GB 2001-4534	20010223
				GB 2001-4534	20010223

OS MARPAT 138:73178
 IT 328083-09-6P, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[3-(2-pyridinyl)-1,2,4-oxadiazol-5-yl]acetamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (CCR3 antagonist; prepn. and pharmaceutical combinations of
 [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3
 antagonists for treatment of asthma, allergic disease, or inflammation)
 RN 328083-09-6 CAPLUS
 CN 1,2,4-Oxadiazole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-(2-pyridinyl)- (9CI) (CA INDEX NAME)



GI



AB Title compds. I [wherein Z = CR₄R₅, CO, or CR₄R₅Z₁; Z₁ = alkylene, alkenylene, or CONH; R₁ = (un)substituted alkyl, alkenyl, (hetero)cycloalkyl, or (hetero)aryl; Q = O, S, NR₉, CO, CONR₉, NR₉CO, or CH=CH; m = 0-1; n = 0-6 with the proviso that when n = 0; then m = 0; R₂ and R₃ = independently H or alkyl; or CR₂R₃ = (alkyl)cycloalkyl; T = NR₁₀, CONR₁₀, NR₁₁CONR₁₀, or CONR₁₀R₁₁; X₁-X₄ = independently CH₂CHR₁₂ or CO; R₄ and R₅ = independently H or alkyl; R₆ = (un)substituted (hetero)aryl; R₉-R₁₁ = independently H, alkyl, haloalkyl, hydroxyalkyl, cycloalkyl(alkyl), or phenylalkyl; R₁₂ = independently (cyclo)alkyl or CO; or R₁₂ groups of X₁ and X₃ or X₄, or X₂ and X₃ or X₄ join to form CH₂CH₂, CH₂CH₂CH₂, CH₂OCH₂, or CH₂SCH₂; or pharmaceutically acceptable salts or solvates thereof] were prep'd. as cysteine-cysteine chemokine receptor 3 (CCR3) antagonists for use in pharmaceutical combinations with a histamine antagonist, steroid, leukotriene modulator, human cytokine, .beta.-agonist, phosphodiesterase inhibitor, or antibody (no data). For example, 1-(3,4-dichlorobenzyl)-4-piperidinamine.bul.2CF₃CO₂H was condensed with 2-(4-fluorophenyl)acetic acid to give N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]-2-(4-fluorophenyl)acetamide (II). I are useful in combination therapy for the treatment of asthma, rhinitis, and other allergic or inflammatory conditions (no data).

L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2001:152644 CAPLUS

DN 134:207822

TI Preparation of substituted piperidines as modulators of chemokine receptor activity

IN Thom, Stephen; Baxter, Andrew; Kindon, Nicholas; McInally, Thomas; Springthorpe, Brian; Perry, Matthew; Harden, David; Evans, Richard; Marriott, David

PA AstraZeneca UK Limited, UK

SO PCT Int. Appl., 133 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001014333	A1	20010301	WO 2000-GB3179	20000818
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA,				

ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
 CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 SE 1999-2987 A 19990824

EP 1212299 A1 20020612 EP 2000-951768 20000818
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL
 SE 1999-2987 A 19990824

WO 2000-GB3179 W 20000818
 JP 2003507456 T2 20030225 JP 2001-518423 20000818
 SE 1999-2987 A 19990824
 WO 2000-GB3179 W 20000818


OS MARPAT 134:207822

IT 328083-09-6P



RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of substituted piperidines as modulators of chemokine receptor activity)



RN 328083-09-6 CAPLUS

CN 1,2,4-Oxadiazole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-(2-pyridinyl)- (9CI) (CA INDEX NAME)



GI





AB The title compds. [I; Z = CR₄R₅, CO, CR₄R₅Z₁; Z₁ = alkylene, alkenylene, CONH; R₁ = (un)substituted alkyl, alkenyl, 3-14 membered (un)satd. ring system which optionally further comprises up to two ring carbon atoms that form carbonyl groups and which optionally further comprises up to 4 ring heteroatoms selected from N, O, and S; m = 0-1; Q = O, S, CO, etc.; n = 0-6 (when n = 0, then m = 0); R₂, R₃ = H, alkyl; (CR₂R₃)_n = cycloalkyl optionally substituted by alkyl; T = NR₁₀, CONR₁₀, NR₁₁CONR₁₀, etc.; X₁-X₄ = CH₂, CHR₁₂ (wherein R₁₂ = alkyl, cycloalkyl(alkyl), CO, etc.); R₄, R₅ = H, alkyl; R₆ = (un)substituted aryl, heterocyclyl; R₁₀-R₁₁ = H, alkyl, haloalkyl, etc.] and their pharmaceutically acceptable salts, useful in therapy, esp. for the treatment of chemokine receptor related diseases (such as inflammatory disease) and conditions, were prep'd. E.g., a 3-step synthesis of the piperidine II was given. The exemplified compds. I were found to be antagonists of the eotaxin mediated [Ca²⁺]_i in human eosinophils and/or antagonists of the MIP-1. α . mediated [Ca²⁺]_i in human monocytes (no data). Certain compds. I were found to be antagonists of the eotaxin mediated human eosinophil chemotaxis (no data).

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s chemikine receptor and CCR1 and CCR3
L9 0 CHEMIKINE RECEPTOR AND CCR1 AND CCR3

=> s chemokine reprot and G-protein
L10 0 CHEMOKINE REPTOR AND G-PROTEIN

=> s piperidine and chemokine receptor
L11 145 PIPERIDINE AND CHEMOKINE RECEPTOR

=> s l11 and CCR1 and CCR3
L12 4 L11 AND CCR1 AND CCR3

=> d l12 fbib hitstr abs total



L12 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2002:617868 CAPLUS

TI Discovery and structure-activity relationship of N-(ureidoalkyl)-benzyl-piperidines as potent small molecule CC chemokine receptor-3 antagonists
AU De Lucca, George V.; Kim, Ui T.; Vargo, Brian; Welch, Patricia K.; Johnson, Curt; Covington, Maryanne; Davies, Paul; Solomon, Kimberly A.; Newton, Robert C.; Trainor, George L.; Decicco, Carl P.; Ko, Soo S.
CS Medicinal Chemistry, Bristol-Myers Squibb, Wilmington, DE, 19880-0336, USA
SO Abstracts of Papers, 224th ACS National Meeting, Boston, MA, United States, August 18-22, 2002 (2002), MEDI-014 Publisher: American Chemical Society, Washington, D. C.
CODEN: 69CZPZ
DT Conference; Meeting Abstract
LA English
AB Eosinophils have been assocd. with disease symptoms in allergic asthma. Eotaxin, a member of the CC chemokine family, is the major chemokine responsible for eosinophil recruitment and activation into the lungs of asthmatic patients. The eotaxin receptor has been identified as CCR3. Thus, CCR3 receptor antagonists may be useful for the treatment of allergic asthma. Starting with several initial hits from screening our inhouse library of compds., we were able to define essential pharmacophore features that are responsible for CCR3 binding affinity. Further structure-activity studies lead to a series of compds. that were potent and specific CCR3 receptor antagonist with IC50 less than 10 nM. They are also greater than 100 fold selective over other chemokine receptors (CCR1, CCR2, CCR5) and other seven transmembrane receptors. The SAR, oral bioavailability, and the functional activity of this series of compds. will be discussed and presented.

L12 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2002:515125 CAPLUS
DN 137:210415
TI Discovery and structure-activity relationship of N-(ureidoalkyl)-benzyl-piperidines as potent small molecule CC chemokine receptor-3 (CCR3) antagonists
AU De Lucca, George V.; Kim, Ui T.; Johnson, Curt; Vargo, Brian J.; Welch, Patricia K.; Covington, Maryanne; Davies, Paul; Solomon, Kimberly A.; Newton, Robert C.; Trainor, George L.; Decicco, Carl P.; Ko, Soo S.
CS Experimental Station, Bristol-Myers Squibb Company, Wilmington, DE, 19880-0336, USA
SO Journal of Medicinal Chemistry (2002), 45(17), 3794-3804
CODEN: JMCMAR; ISSN: 0022-2623
PB American Chemical Society
DT Journal
LA English
AB Structure-activity relationship (SAR) studies of initial screening hits from our corporate library of compds. and a structurally related series of CCR1 receptor antagonists were used to det. that an N-(alkyl)benzylpiperidine is an essential pharmacophore for selective CCR3 antagonists. Further SAR studies that introduced N-(ureidoalkyl) substituents improved the binding potency of these compds. from the micromolar to the low nanomolar range. This new series of compds. also displays highly potent, in vitro functional CCR3 -mediated antagonism of eotaxin-induced Ca²⁺ mobilization and chemotaxis of human eosinophils.

RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2001:321173 CAPLUS
 DN 135:162081
 TI Discovery of a novel CCR3 selective antagonist
 AU Naya, A.; Kobayashi, K.; Ishikawa, M.; Ohwaki, K.; Saeki, T.; Noguchi, K.; Ohtake, N.
 CS Banyu Tsukuba Research Institute, Tsukuba, Ibaraki, 300-2611, Japan
 SO Bioorganic & Medicinal Chemistry Letters (2001), 11(9), 1219-1223
 CODEN: BMCLE8; ISSN: 0960-894X
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 OS CASREACT 135:162081
 GI



AB In searching for a novel CCR3 receptor antagonist, we designed a library that included a variety of carboxamide derivs. based on the structure of our potent antagonists for human CCR1 and CCR3 receptors, and screened the new compds. for inhibitory activity against ^{125}I -Eotaxin binding to human CCR3 receptors expressed in CHO cells. Among them, two 2-(benzothiazolethio)acetamide derivs. (I and II) showed binding affinities with IC₅₀ values of 750 and 1000 nM, resp., for human CCR3 receptors. I and II also possessed weak binding affinities for human CCR1 receptors. We selected I as a lead compd. for derivatization to improve in vitro potency and selectivity for CCR3 over CCR1 receptors. Derivatization of I by incorporating substituents into each benzene ring of the benzothiazole and piperidine side chain resulted in the discovery of a compd. (III) exhibiting 820-fold selectivity for CCR3 receptors ($\text{IC}_{50}=2.3$ nM) over CCR1 receptors ($\text{IC}_{50}=1900$ nM). III also showed potent functional antagonist activity for

inhibiting Eotaxin (IC₅₀=27 nM) - or RANTES (IC₅₀=13 nM) -induced Ca²⁺ increases in eosinophils.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2000:603235 CAPLUS
DN 133:290649
TI Identification of the binding site for a novel class of CCR2b
chemokine receptor antagonists: binding to a common
chemokine receptor motif within the helical bundle
AU Mirzadegan, Tara; Diehl, Frank; Ebi, Bettina; Bhakta, Sunil; Polksy,
Irene; McCarley, Deborah; Mulkins, Mary; Weatherhead, Gabe S.; Lapierre,
Jean-Marc; Dankwardt, John; Morgans, David, Jr.; Wilhelm, Robert;
Jarnagin, Kurt
CS Roche Bioscience, Palo Alto, CA, 94304, USA
SO Journal of Biological Chemistry (2000), 275(33), 25562-25571
CODEN: JBCHA3; ISSN: 0021-9258
PB American Society for Biochemistry and Molecular Biology
DT Journal
LA English
AB Monocyte chemoattractant-1 (MCP-1) stimulates leukocyte chemotaxis to inflammatory sites, such as rheumatoid arthritis, atherosclerosis, and asthma, by use of the MCP-1 receptor, CCR2, a member of the G-protein-coupled seven-transmembrane receptor superfamily. These studies identified a family of antagonists, spiropiperidines. One of the more potent compds. blocks MCP-1 binding to CCR2 with a Kd of 60 nM, but it is unable to block binding to CXCR1, CCR1, or CCR3. These compds. were effective inhibitors of chemotaxis toward MCP-1 but were very poor inhibitors of CCR1-mediated chemotaxis. The compds. are effective blockers of MCP-1-driven inhibition of adenylate cyclase and MCP-1- and MCP-3-driven cytosolic calcium influx; the compds. are not agonists for these pathways. The authors showed that glutamate 291 (Glu291) of CCR2 is a crit. residue for high affinity binding and that this residue contributes little to MCP-1 binding to CCR2. The basic nitrogen present in the spiropiperidine compds. may be the interaction partner for Glu291, because the basicity of this nitrogen was essential for affinity; furthermore, a different class of antagonists, a class that does not have a basic nitrogen (2-carboxypyrrroles), were not affected by mutations of Glu291. In addn. to the CCR2 receptor, spiropiperidine compds. have affinity for several biogenic amine receptors. Receptor models indicate that the acidic residue, Glu291, from transmembrane-7 of CCR2 is in a position similar to the acidic residue contributed from transmembrane-3 of biogenic amine receptors, which may account for the shared affinity of spiropiperidines for these two receptor classes. The models suggest that the acid-base pair, Glu291 to piperidine nitrogen, anchors the spiropiperidine compd. within the transmembrane ovoid bundle. This binding site may overlap with the space required by MCP-1 during binding and signaling; thus the small mol. ligands act as antagonists. An acidic residue in transmembrane region 7 is found in most chemokine receptors and is rare in other serpentine receptors. The model of the binding site may suggest ways to make new small mol. chemokine receptor antagonists, and it may rationalize the design of more potent and selective antagonists.

RE.CNT 52 THERE ARE 52 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

10069215.1

Page 16

=> log y		
COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	44.04	345.91
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-3.91	-3.91

STN INTERNATIONAL LOGOFF AT 10:47:53 ON 13 OCT 2003

Welcome to STN International! Enter x:x

LOGINID: ssspta1611sxp

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 10:51:12 ON 13 OCT 2003

FILE 'REGISTRY' ENTERED AT 10:52:17 ON 13 OCT 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 OCT 2003 HIGHEST RN 603065-76-5

DICTIONARY FILE UPDATES: 12 OCT 2003 HIGHEST RN 603065-76-5

TSCA INFORMATION NOW CURRENT THROUGH JULY 14, 2003

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:

<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

=>


Uploading 10069215.2

L1 STRUCTURE UPLOADED

=> d 11

L1 HAS NO ANSWERS

L1 STR



G1 O,N,C,S,CH ,

G2 Cb,Cy,Hy

Structure attributes must be viewed using STN Express query preparation.

```
=> s 11 sss full
FULL SEARCH INITIATED 10:52:43 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 64899 TO ITERATE

100.0% PROCESSED 64899 ITERATIONS          4255 ANSWERS
SEARCH TIME: 00.00.02
```

L2 4255 SEA SSS FUL L1

```
=> file caplus
COST IN U.S. DOLLARS           SINCE FILE      TOTAL
                                ENTRY          SESSION
FULL ESTIMATED COST          148.55         148.97
```

```
FILE 'CAPLUS' ENTERED AT 10:53:24 ON 13 OCT 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)
```

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 13 Oct 2003 VOL 139 ISS 16
FILE LAST UPDATED: 12 Oct 2003 (20031012/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> s 12
L3      1572 L2

=> s 13 and oxadiazole
L4      13 L3 AND OXADIAZOLE

=> s 13 and imidazole
L5      98 L3 AND IMIDAZOLE

=> s 13 and oxazole
L6      6 L3 AND OXAZOLE

=> s 13 and one
L7      295 L3 AND ONE

=> s 13 and chemikine
L8      0 L3 AND CHEMIKINE
```

```
=> s l3 and chemokine and ccr1 and ccr3
L9          1 L3 AND CHEMOKINE AND CCR1 AND CCR3

=> d his

      (FILE 'HOME' ENTERED AT 10:51:12 ON 13 OCT 2003)

      FILE 'REGISTRY' ENTERED AT 10:52:17 ON 13 OCT 2003
L1          STRUCTURE UPLOADED
L2          4255 S L1 SSS FULL

      FILE 'CAPLUS' ENTERED AT 10:53:24 ON 13 OCT 2003
L3          1572 S L2
L4          13 S L3 AND OXADIAZOLE
L5          98 S L3 AND IMIDAZOLE
L6          6 S L3 AND OXAZOLE
L7          295 S L3 AND ONE
L8          0 S L3 AND CHEMIKINE
L9          1 S L3 AND CHEMOKINE AND CCR1 AND CCR3

=> s l3 and triazole
L10         20 L3 AND TRIAZOLE

=> s l3 and 1,2,4-triazole
L11         15 L3 AND 1,2,4-TRIAZOLE

=> d l4 fbib hitstr abs totasl
'TOTASL' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'
```

The following are valid formats:

```
ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data and PI table (default)
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing
FAM ----- AN, PI and PRAI in table, plus Patent Family data
FBIB ----- AN, BIB, plus Patent FAM
IND ----- Indexing data
IPC ----- International Patent Classifications
MAX ----- ALL, plus Patent FAM, RE
PAT----- PI, SO
SAM ----- CC, SX, TI, ST, IT
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
                      SCAN must be entered on the same line as the DISPLAY,
                      e.g., D SCAN or DISPLAY SCAN)
STD ----- BIB, IPC, and NCL

IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IMAX ----- MAX, indented with text labels
ISTD ----- STD, indented with text labels

OBIB ----- AN, plus Bibliographic Data (original)
```

OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
 SIBIB ----- IBIB, no citations

HIT ----- Fields containing hit terms
 HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT) containing hit terms
 HITRN ----- HIT RN and its text modification
 HITSTR ----- HIT RN, its text modification, its CA index name, and its structure diagram
 HITSEQ ----- HIT RN, its text modification, its CA index name, its structure diagram, plus NTE and SEQ fields
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and its structure diagram
 FHITSEQ ----- First HIT RN, its text modification, its CA index name, its structure diagram, plus NTE and SEQ fields
 KWIC ----- Hit term plus 20 words on either side
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.

ENTER DISPLAY FORMAT (BIB):BIB

L4 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:610204 CAPLUS
 DN 139:164801
 TI Preparation of 2,4-pyrimidinediamines as IgE and/or IgG receptor modulators for treatment of allergic diseases, inflammatory conditions, and tissue destruction
 IN Singh, Rajinder; Argade, Ankush; Payan, Donald G.; Molineaux, Susan; Holland, Sacha J.; Clough, Jeffrey; Keim, Holger; Bhamidipati, Somasekhar; Sylvain, Catherine; Li, Weigun; Rossi, Alexander B.
 PA Rigel Pharmaceuticals, Inc., USA
 SO PCT Int. Appl., 648 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003063794	A2	20030807	WO 2003-US3022	20030131
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,

CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG
PRAI US 2002-353267P P 20020201
US 2002-353333P P 20020201
US 2002-399673P P 20020729
US 2002-434277P P 20021217
OS MARPAT 139:164801

=> d his

(FILE 'HOME' ENTERED AT 10:51:12 ON 13 OCT 2003)

FILE 'REGISTRY' ENTERED AT 10:52:17 ON 13 OCT 2003

L1 STRUCTURE uploaded
L2 4255 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 10:53:24 ON 13 OCT 2003

L3 1572 S L2
L4 13 S L3 AND OXADIAZOLE
L5 98 S L3 AND IMIDAZOLE
L6 6 S L3 AND OXAZOLE
L7 295 S L3 AND ONE
L8 0 S L3 AND CHEMIKINE
L9 1 S L3 AND CHEMOKINE AND CCR1 AND CCR3
L10 20 S L3 AND TRIAZOLE
L11 15 S L3 AND 1,2,4-TRIAZOLE

=> d 14 fbib hitstr abs total

L4 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2003:610204 CAPLUS
DN 139:164801

TI Preparation of 2,4-pyrimidinediamines as IgE and/or IgG receptor modulators for treatment of allergic diseases, inflammatory conditions, and tissue destruction

IN Singh, Rajinder; Argade, Ankush; Payan, Donald G.; Molineaux, Susan; Holland, Sacha J.; Clough, Jeffrey; Keim, Holger; Bhamidipati, Somasekhar; Sylvain, Catherine; Li, Weigun; Rossi, Alexander B.

PA Rigel Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 648 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003063794	A2	20030807	WO 2003-US3022	20030131
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,

NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG

US 2002-353267PP 20020201
US 2002-353333PP 20020201
US 2002-399673PP 20020729
US 2002-434277PP 20021217

OS MARPAT 139:164801


IT 575479-15-1P 575479-16-2P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(IgE and/or IgG receptor modulator; prepn. of pyrimidinediamines as IgE and/or IgG receptor modulators for treatment of allergic diseases, inflammatory conditions, and tissue destruction)


RN 575479-15-1 CAPPLUS

CN 2,4-Pyrimidinediamine, N4-(3,4-dimethoxyphenyl)-5-fluoro-N2-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)





RN 575479-16-2 CAPPLUS

CN Phenol, 3-[[5-fluoro-2-[[1-(phenylmethyl)-4-piperidinyl]amino]-4-pyrimidinyl]amino]- (9CI) (CA INDEX NAME)



GI



AB Title compds. I [wherein L1 and L2 = independently a bond or a linker; R2 = (un)substituted alkyl, (hetero)cycloalkyl, or (hetero)aryl; R4 = H or R2; R5 = R6 or (un)substituted alkyl, alkenyl, or alkynyl; R6 = independently H, an electroneg. group, protected alc. or thiol, haloalkyl(oxy), halo, CN, NC, OCN, SCN, NO, NO₂, N₃, or (un)substituted amino, sulfamoyl(oxy), acyl, carboxy, carbamoyl, (hetero)aryl(alkyl), etc.; with provisos and exclusions; and salts, hydrates, solvates, N-oxides, and prodrugs thereof] were prep'd. as inhibitors of the IgE and/or IgG receptor signaling cascades that lead to the release of chem. mediators. For example, coupling of 2,4-dichloropyrimidine with 4-ethoxyaniline in EtOH provided N₂,N₄-bis(4-ethoxyphenyl)-2,4-pyrimidinediamine (II). The latter inhibited degranulation of bone marrow derived mast cells challenged with anti-IgE and ionomycin with IC₅₀ values of 4.5 .mu.M and 4.4 .mu.M, resp. Thus, I and their pharmaceutical compns. are useful in the treatment and prevention of diseases characterized by, caused by, or assocd. with the release of chem. mediators via degranulation of mast, basophil, neutrophil, or eosinophil cells and other processes effected by activation of the IgE and/or IgG receptor signaling cascades. The treatment and prevention of allergic diseases, low grade scarring, diseases assocd. with tissue destruction, diseases assocd. with tissue inflammation, inflammation, and scarring are targeted uses (no data).

L4 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:319886 CAPLUS

DN 138:338155

TI Preparation of oxadiazolyl-biphenylcarboxamides as p38 kinase inhibitors

IN Angell, Richard Martyn; Bamborough, Paul; Cockerill, George Stuart; Smith, Kathryn Jane; Walker, Ann Louise

PA Glaxo Group Limited, UK

SO PCT Int. Appl., 44 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.


KIND DATE

APPLICATION NO. DATE

PI WO 2003033482 A1 20030424 WO 2002-EP11574 20021016
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD,
 RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
 PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
 NE, SN, TD, TG

GB 2001-24932 A 20011017

OS MARPAT 138:338155
 IT 515143-78-9P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of oxadiazolyl-biphenylcarboxamides as p38 kinase inhibitors)
 RN 515143-78-9 CAPLUS
 CN [1,1'-Biphenyl]-4-carboxamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]-2'-methyl-5'-(5-methyl-1,3,4-oxadiazol-2-yl) - (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title compds. [I; X = a bond, (un)substituted Ph; R1 = (un)substituted 5-7 membered heterocyclyl, 5-7 membered heteroaryl, fused bicycyl; R2 = H, alkyl, (CH₂)pcycloalkyl; or when X = a bond and m and n are both zero, NR1R2 = 5-6 membered heterocyclyl optionally contg. one addnl. heteroatom selected from O and N which can be optionally substituted by alkyl; R3 = II (wherein R4 = H, alkyl); U = Me, halo; V, Y = H, Me, halo; m, n = 0-2; m + n = 0-4; p = 0-1; r = 0-2; with the provisos], useful as pharmaceuticals, particularly as p38 kinase inhibitors, were prep'd. E.g., a 6-step synthesis of the carboxamide III, starting from 3-bromo-4-methylbenzoic acid, was given.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN


AN 2003:44146 CAPLUS
 DN 138:73178
 TI Preparation and pharmaceutical combinations of
 [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3 antagonists
 for treatment of asthma, allergic disease, or inflammation
 IN Bahl, Ash; Perry, Matthew; Springthorpe, Brian
 PA AstraZeneca AB, Swed.
 SO Brit. UK Pat. Appl., 91 pp.
 CODEN: BAXXDU
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 2373186	A1	20020918	GB 2001-4534	20010223
				GB 2001-4534	20010223


OS MARPAT 138:73178
 IT 328082-81-1P, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-methyl-
 1-phenyl-1H-pyrazol-4-yl)acetamide 328082-83-3P,
 2-[2-(4-Chlorophenyl)-5-methyl-1,3-thiazol-4-yl]-N-[1-(3,4-dichlorobenzyl)-
 4-piperidinyl]acetamide 328082-85-5P, N-[1-(3,4-Dichlorobenzyl)-
 4-piperidinyl]-2-(4-fluorophenyl)acetamide 328082-86-6P,
 N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[2-(2-pyrazinyl)-1,3-thiazol-4-
 yl]acetamide 328082-89-9P, N-[1-(3,4-Dichlorobenzyl)-4-
 piperidinyl]-1H-benzimidazol-2-amine 328082-91-3P,
 N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N'-(3,4-dichlorophenyl)urea
 328082-92-4P, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N'-(3-
 methoxyphenyl)urea 328082-93-5P, N-[1-(3,4-Dichlorobenzyl)-4-
 piperidinyl]-N-(4-methoxybenzyl)amine dihydrochloride 328083-00-7P
 , Cis-N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[3-(2-pyridinyl)-1,2,4-
 oxadiazol-5-yl]cyclopropanecarboxamide 328083-04-1P,
 N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-phenyl-1H-1,2,4-triazol-5-
 yl)acetamide 328083-07-4P, N-[1-(3,4-Dichlorobenzyl)-4-
 piperidinyl]-2-(5-phenyl-1,3,4-oxadiazol-2-yl)acetamide acetate
 328083-09-6P, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[3-(2-
 pyridinyl)-1,2,4-oxadiazol-5-yl]acetamide 328083-11-0P,
 N-[1-(4-Bromobenzyl)-4-piperidinyl]-2-(4-fluorophenyl)acetamide
 328083-36-9P, N-[1-[(3,4-Dichlorophenyl)methyl]-4-piperidinyl]-4-
 hydroxybenzeneacetamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)
 (CCR3 antagonist; prepn. and pharmaceutical combinations of
 [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3
 antagonists for treatment of asthma, allergic disease, or inflammation)

RN 328082-81-1 CAPLUS
 CN 1H-Pyrazole-4-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-
 5-methyl-1-phenyl- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




RN 328082-83-3 CAPLUS

CN 4-Thiazoleacetamide, 2-(4-chlorophenyl)-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methyl- (9CI) (CA INDEX NAME)




RN 328082-85-5 CAPLUS


CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)




RN 328082-86-6 CAPLUS
 CN 4-Thiazoleacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-pyrazinyl- (9CI) (CA INDEX NAME)




RN 328082-89-9 CAPLUS
 CN 1H-Benzimidazol-2-amine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-91-3 CAPLUS
 CN Urea, N-(3,4-dichlorophenyl)-N'-(1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl)- (9CI) (CA INDEX NAME)



RN 328082-92-4 CAPLUS
 CN Urea, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-N'-(3-methoxyphenyl)- (9CI) (CA INDEX NAME)




RN 328082-93-5 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-methoxyphenyl)methyl]-, dihydrochloride (9CI) (CA INDEX NAME)




● 2 HCl

RN 328083-00-7 CAPLUS
 CN Cyclopropanecarboxamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-[3-(2-pyridinyl)-1,2,4-oxadiazol-5-yl]-, (1R,2S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 328083-04-1 CAPLUS
 CN 1H-1,2,4-Triazole-3-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-phenyl- (9CI) (CA INDEX NAME)




RN 328083-07-4 CAPLUS

CN 1,3,4-Oxadiazole-2-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-phenyl-, monoacetate (9CI) (CA INDEX NAME)


CM 1

CRN 328083-06-3
CMF C22 H22 Cl2 N4 O2




CM 2

CRN 64-19-7
CMF C2 H4 O2




RN 328083-09-6 CAPLUS

CN 1,2,4-Oxadiazole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-(2-pyridinyl)- (9CI) (CA INDEX NAME)



RN 328083-11-0 CAPLUS


CN Benzeneacetamide, N-[1-[(4-bromophenyl)methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)



RN 328083-36-9 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-

hydroxy- (9CI) (CA INDEX NAME)



IT 328081-86-3, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-methylbenzyl)amine 328081-87-4, N-[4-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]phenyl]acetamide 328081-88-5, 3-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]phenol 328081-89-6, N-[(4-Chloro-1-methyl-1H-pyrazol-3-yl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine 328081-90-9, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(5-methyl-2-furyl)methyl]amine 328081-91-0, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-nitrobenzyl)amine 328081-92-1, N-Benzyl-1-(3,4-dichlorobenzyl)-4-piperidinamine 328081-93-2, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-fluorobenzyl)amine 328081-94-3, N-(2,6-Dichlorobenzyl)-1-(3,4-dichlorobenzyl)-4-piperidinamine 328081-95-4, N,1-Bis(3,4-dichlorobenzyl)-4-piperidinamine 328081-96-5, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(2-pyridinylmethyl)amine 328081-97-6, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(3-methyl-2-thienyl)methyl]amine 328081-98-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(5-methyl-2-thienyl)methyl]amine 328081-99-8, 5-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-methoxyphenol 328082-00-4, 4-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-nitrophenol 328082-01-5, 3-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-4H-chromen-4-one 328082-02-6, N-[(5-Chloro-1,3-dimethyl-1H-pyrazol-4-yl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine 328082-03-7, N-[(4-Chloro-1H-pyrazol-3-yl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine 328082-04-8, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[[1-(4-methylbenzyl)-1H-pyrazol-5-yl]methyl]amine 328082-05-9, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(2-phenyl-1H-imidazol-4-yl)methyl]amine 328082-06-0, N-[(2-Chloro-3-quinolinyl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine 328082-08-2, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(3-quinolinylmethyl)amine 328082-09-3, [5-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-furyl]methyl acetate 328082-10-6, 4-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-1,5-dimethyl-2-phenyl-1,2-dihydro-3H-pyrazol-3-one 328082-11-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-pyridinylmethyl)amine 328082-12-8, 5-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-nitrophenol 328082-13-9, N-[2-(tert-Butylsulfanyl)benzyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine 328082-14-0, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-ethylbenzyl)amine 328082-15-1, 5-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-hydroxybenzoic acid 328082-16-2, N-(1,3-Benzodioxol-4-ylmethyl)-1-(3,4-dichlorobenzyl)-4-piperidinamine 328082-17-3, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(1,3-thiazol-2-ylmethyl)amine 328082-18-4, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(5-ethyl-2-furyl)methyl]amine 328082-19-5, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(2-quinolinylmethyl)amine 328082-20-8, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-

quinolinylmethyl)amine **328082-21-9**, 5-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-hydroxy-3-methoxybenzoic acid
328082-22-0, N-[(4-Bromo-1H-pyrazol-3-yl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine **328082-23-1**,
2-[2-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-6-methoxyphenoxy]acetic acid **328082-24-2**, N-[(4-Bromo-1-methyl-1H-pyrazol-3-yl)methyl]-1-(3,4-dichlorobenzyl)-4-piperidinamine
328082-25-3, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-iodobenzyl)amine **328082-26-4**, 3-[[[1-(3,4-Dichlorobenzyl)-4-piperidinyl]amino]methyl]-6,7-dimethyl-4H-chromen-4-one
328082-27-5, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(4-isopropoxybenzyl)amine **328082-28-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(1-methyl-1H-benzimidazol-2-yl)methyl]amine
328082-29-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(3-methylbenzyl)amine **328082-30-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(3-pyridinylmethyl)amine **328082-31-1**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-(2,4-dimethylbenzyl)amine
328082-32-2, Ethyl 5-[[[1-(3,4-dichlorobenzyl)-4-piperidinyl]amino]methyl]-2-methyl-3-furoate **328082-33-3**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-3-furamide **328082-36-6**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-6-methoxy-4-quinolinecarboxamide
328082-37-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-furyl)-4-quinolinecarboxamide **328082-40-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,5-dimethoxyphenyl)acetamide **328082-41-3**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-methoxyphenyl)acetamide
328082-42-4, 2-(5-Chloro-2-oxo-1,3-benzothiazol-3(2H)-yl)-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]acetamide **328082-44-6**,
2-(Benzothiophen-3-yl)-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]acetamide
328082-47-9, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-3-fluoro-2-methylbenzamide **328082-48-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N'-(1-phenylethyl)phthalamide **328082-49-1**,
2-Cyclopentyl-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]acetamide
328082-50-4, 4-Chloro-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]-2-nitrobenzamide **328082-51-5**, 2,2-Dichloro-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]-1-methylcyclopropanecarboxamide
328082-52-6, tert-Butyl 4-[5-[[[1-(3,4-dichlorobenzyl)-4-piperidinyl]amino]carbonyl]-2-methoxyphenyl]-1-piperazinecarboxylate
328082-53-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-5-oxo-1-(2-thienylmethyl)-3-pyrrolidinecarboxamide **328082-55-9**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-4-fluorobenzamide
328082-56-0, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-methylbenzamide **328082-57-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-3-methylbenzamide **328082-58-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-4-(hydroxymethyl)benzamide
328083-64-3, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)acetamide **328083-73-4**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[4-(methylsulfonyl)benzyl]amine
328083-75-6, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-phenyl-1,3-thiazol-2-yl)acetamide **328083-76-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-phenyl-1,3-thiazol-4-yl)acetamide
389062-07-1, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-phenylacetamide **479554-54-6**, 2-([1,1'-Biphenyl]-4-yloxy)-N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]acetamide **479554-59-1**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-bromophenyl)acetamide
479554-60-4, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-aminophenyl)acetamide **479554-61-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-bromophenyl)acetamide **479554-62-6**,
N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-methylphenyl)acetamide

479554-63-7, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-methylphenyl)acetamide **479554-64-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-chloro-4-hydroxyphenyl)acetamide **479554-65-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-nitrophenyl)acetamide **479554-66-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-chlorophenyl)acetamide **479554-67-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-chlorophenyl)acetamide **479554-69-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-nitrophenyl)acetamide **479554-71-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,4-dimethoxyphenyl)acetamide **479554-72-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-fluoro-4-hydroxyphenyl)acetamide **479554-73-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,4-methylenedioxyphenyl)acetamide **479554-75-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-phenylphenyl)acetamide **479554-76-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,4-dichlorophenyl)acetamide **479554-82-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-methylphenyl)acetamide **479554-83-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[4-(trifluoromethoxy)phenyl]acetamide **479554-84-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-methoxyphenyl)acetamide **479554-85-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[4-(dimethylamino)phenyl]acetamide **479554-87-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,4,5-trimethoxyphenyl)acetamide **479554-89-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-aminophenyl)acetamide **479554-90-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(1-naphthyl)acetamide **479554-91-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-methoxy-4-hydroxyphenyl)acetamide **479554-92-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[3-[[6-bromo-1-(prop-2-en-1-yl)-2-naphthyl]oxy]methyl]phenyl]acetamide **479554-93-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-[4-(4-nitrobenzyloxy)phenyl]acetamide **479554-94-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-fluoro-4-methoxyphenyl)acetamide **479554-96-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-hydroxyphenyl)acetamide **479554-97-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-(benzyloxy)phenyl)acetamide **479554-98-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-(3-nitrophenyl)phenyl)acetamide **479554-99-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,5-dimethylphenyl)acetamide **479555-00-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-iodophenyl)acetamide **479555-01-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-bromophenyl)acetamide **479555-02-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-methyl-3-nitrophenyl)acetamide **479555-03-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-hydroxy-4-methoxyphenyl)acetamide **479555-04-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-fluorophenyl)acetamide **479555-05-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-fluorophenyl)acetamide **479555-06-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-chlorophenyl)acetamide **479555-07-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,5-dimethylphenyl)acetamide **479555-09-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,4-difluorophenyl)acetamide **479555-10-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,4-difluorophenyl)acetamide **479555-11-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,5-difluorophenyl)acetamide **479555-12-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-pyridinyl)acetamide **479555-13-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-pyridinyl)acetamide **479555-14-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-bromo-3-pyridinyl)acetamide

479555-15-2, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,4-dimethoxyphenyl)acetamide **479555-16-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-(benzyloxy)phenyl)acetamide **479555-17-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-methyl-1-naphthyl)acetamide **479555-18-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-ethoxyphenyl)acetamide **479555-19-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-butoxyphenyl)acetamide **479555-20-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(1-indolyl)acetamide **479555-21-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-thienyl)acetamide **479555-22-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,4-dichlorophenyl)acetamide **479555-23-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,6-dichlorophenyl)acetamide **479555-24-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-(2-chloroacetamido)-4-thiazolyl)acetamide **479555-25-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,5-dimethoxyphenyl)acetamide **479555-26-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-indolyl)acetamide **479555-27-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-methoxy-3-indolyl)acetamide **479555-28-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-naphthyl)acetamide **479555-29-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-(methylsulfonyl)phenyl)acetamide **479555-30-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,4,6-trimethylphenyl)acetamide **479555-31-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-isopropyl-2-methyl-3-indolyl)acetamide **479555-32-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-(1-pyrrolidinyl)-2H-tetrazol-2-yl)acetamide **479555-33-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-(4-methylphenyl)-2H-tetrazol-2-yl)acetamide **479555-34-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-methoxyphenyl)acetamide **479555-35-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-chloro-3-benzothienyl)acetamide **479555-36-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-methyl-2-phenyl-4-thiazolyl)acetamide **479555-37-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-chloro-3-methyl-2-benzothienyl)acetamide **479555-38-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-methyl-2-benzothienyl)acetamide **479555-39-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3-nitro-1,2,4-triazol-1-yl)acetamide **479555-40-3**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(5-methyl-3,4-dinitro-1-pyrazolyl)acetamide **479555-41-4**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-(3-methylbutoxy)phenyl)acetamide **479555-42-5**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,3-dimethyl-5-indolyl)acetamide **479555-43-6**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-chloro-3,5-dimethyl-1-pyrazolyl)acetamide **479555-44-7**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-nitro-3,5-dimethyl-1-pyrazolyl)acetamide **479555-45-8**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2,4-dinitro-1-imidazolyl)acetamide **479555-46-9**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-nitro-1-imidazolyl)acetamide **479555-47-0**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(3,5-dimethyl-1-pyrazolyl)acetamide **479555-48-1**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(4-hexylphenyl)acetamide **479555-49-2**, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-2-(2-cyanophenyl)acetamide **479555-65-2**, N-[1-[(2,5-Dichlorophenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-66-3**, N-[1-[(2,3-Dichlorophenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-67-4**, N-[1-[(4-Fluorophenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-68-5**, N-[1-[(4-Bromo-3-(methoxycarbonyl)phenyl)methyl]-4-piperidinyl]-2-(4-


fluorophenyl)acetamide **479555-69-6**, N-[1-[(4-Nitrophenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-70-9**, N-[1-[(3-Benzoylphenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-74-3**, N-[1-[(4-Methyl-3-nitrophenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-75-4**, N-[1-[(3,4-Dimethylphenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-76-5**, N-[1-[(4-Methoxy-3-methylphenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-77-6**, N-[1-[(4-(2-Carbamoylphenyl)phenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide **479555-78-7**, N-[1-[(4-((2,6-Dichlorophenyl)methyl)sulfonyl)phenyl)methyl]-4-piperidinyl]-2-(4-fluorophenyl)acetamide

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(CCR3 antagonist; prepn. and pharmaceutical combinations of [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3 antagonists for treatment of asthma, allergic disease, or inflammation)


RN 328081-86-3 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-methylphenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-87-4 CAPLUS

CN Acetamide, N-[4-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]phenyl]- (9CI) (CA INDEX NAME)




RN 328081-88-5 CAPLUS


CN Phenol, 3-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]- (9CI) (CA INDEX NAME)




RN 328081-89-6 CAPLUS
 CN 4-Piperidinamine, N-[(4-chloro-1-methyl-1H-pyrazol-3-yl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-90-9 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(5-methyl-2-furanyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-91-0 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-nitrophenyl)methyl]- (9CI) (CA INDEX NAME)



RN 328081-92-1 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 328081-93-2 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-94-3 CAPLUS

CN 4-Piperidinamine, N-[(2,6-dichlorophenyl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-95-4 CAPLUS

CN 4-Piperidinamine, N,1-bis[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-96-5 CAPLUS

CN 2-Pyridinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328081-97-6 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(3-methyl-2-thienyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-98-7 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(5-methyl-2-thienyl)methyl]- (9CI) (CA INDEX NAME)




RN 328081-99-8 CAPLUS

CN Phenol, 5-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-2-methoxy- (9CI) (CA INDEX NAME)




RN 328082-00-4 CAPLUS

CN Phenol, 4-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-2-nitro- (9CI) (CA INDEX NAME)



RN 328082-01-5 CAPLUS


CN 4H-1-Benzopyran-4-one, 3-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]- (9CI) (CA INDEX NAME)



RN 328082-02-6 CAPLUS

CN 4-Piperidinamine, N-[(5-chloro-1,3-dimethyl-1H-pyrazol-4-yl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)

PAGE 1-A





PAGE 2-A



RN 328082-03-7 CAPLUS


CN 4-Piperidinamine, N-[(4-chloro-1H-pyrazol-3-yl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)






RN 328082-04-8 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[[1-[(4-methylphenyl)methyl]-1H-pyrazol-5-yl]methyl]- (9CI) (CA INDEX NAME)




RN 328082-05-9 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(2-phenyl-1H-imidazol-4-yl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-06-0 CAPLUS

CN 3-Quinolinemethanamine, 2-chloro-N-[(3,4-dichlorophenyl)methyl]-4-piperidinyl- (9CI) (CA INDEX NAME)




RN 328082-08-2 CAPLUS

CN 3-Quinolinemethanamine, N-[(3,4-dichlorophenyl)methyl]-4-piperidinyl- (9CI) (CA INDEX NAME)



RN 328082-09-3 CAPLUS


CN 2-Furanmethanol, 5-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-, acetate (ester) (9CI) (CA INDEX NAME)



RN 328082-10-6 CAPLUS

CN 3H-Pyrazol-3-one, 4-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-1,2-dihydro-1,5-dimethyl-2-phenyl- (9CI) (CA INDEX NAME)


PAGE 1-A




PAGE 2-A




RN 328082-11-7 CAPLUS
 CN 4-Pyridinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl] -
 (9CI) (CA INDEX NAME)




RN 328082-12-8 CAPLUS
 CN Phenol, 5-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-2-nitro- (9CI) (CA INDEX NAME)



RN 328082-13-9 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[[2-[(1,1-dimethylethyl)thio]phenyl]methyl]- (9CI) (CA INDEX NAME)




RN 328082-14-0 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-ethylphenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-15-1 CAPLUS

CN Benzoic acid, 5-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-2-hydroxy- (9CI) (CA INDEX NAME)




RN 328082-16-2 CAPLUS

CN 4-Piperidinamine, N-(1,3-benzodioxol-4-ylmethyl)-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-17-3 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-(2-thiazolylmethyl)- (9CI) (CA INDEX NAME)




RN 328082-18-4 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(5-ethyl-2-furanyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-19-5 CAPLUS

CN 2-Quinolinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-20-8 CAPLUS

CN 4-Quinolinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-21-9 CAPLUS

CN Benzoic acid, 5-[[[1-[(3,4-dichlorophenyl)methyl]amino]methyl]-2-hydroxy-3-methoxy- (9CI) (CA INDEX NAME)




RN 328082-22-0 CAPLUS

CN 4-Piperidinamine, N-[(4-bromo-1H-pyrazol-3-yl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-23-1 CAPLUS


CN Acetic acid, [2-[[[1-[(3,4-dichlorophenyl)methyl]amino]methyl]-6-methoxyphenoxy]- (9CI) (CA INDEX NAME)




RN 328082-24-2 CAPLUS
 CN 4-Piperidinamine, N-[{(4-bromo-1-methyl-1H-pyrazol-3-yl)methyl]-1-[(3,4-dichlorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-25-3 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-iodophenyl)methyl]- (9CI) (CA INDEX NAME)



RN 328082-26-4 CAPLUS
 CN 4H-1-Benzopyran-4-one, 3-[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-6,7-dimethyl- (9CI) (CA INDEX NAME)




RN 328082-27-5 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-(1-methylethoxy)phenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-28-6 CAPLUS

CN 1H-Benzimidazole-2-methanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-1-methyl- (9CI) (CA INDEX NAME)




RN 328082-29-7 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(3-methylphenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-30-0 CAPLUS

CN 3-Pyridinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-31-1 CAPLUS

CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(2,4-dimethylphenyl)methyl]- (9CI) (CA INDEX NAME)




RN 328082-32-2 CAPLUS

CN 3-Furancarboxylic acid, 5-[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]methyl]-2-methyl-, ethyl ester (9CI) (CA INDEX NAME)




RN 328082-33-3 CAPLUS

CN 3-Furancarboxamide, N-[(3,4-dichlorophenyl)methyl]-4-piperidinyl- (9CI) (CA INDEX NAME)




RN 328082-36-6 CAPLUS

CN 4-Quinolinecarboxamide, N-[(3,4-dichlorophenyl)methyl]-4-piperidinyl-6-methoxy- (9CI) (CA INDEX NAME)




RN 328082-37-7 CAPLUS

CN 4-Quinoliniccarboxamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-(2-furanyl)- (9CI) (CA INDEX NAME)




RN 328082-40-2 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-dimethoxy- (9CI) (CA INDEX NAME)




RN 328082-41-3 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methoxy- (9CI) (CA INDEX NAME)




RN 328082-42-4 CAPLUS

CN 3(2H)-Benzothiazoleacetamide, 5-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-oxo- (9CI) (CA INDEX NAME)




RN 328082-44-6 CAPLUS

CN Benzo[b]thiophene-3-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-47-9 CAPLUS


CN Benzamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-fluoro-2-methyl- (9CI) (CA INDEX NAME)




RN 328082-48-0 CAPLUS
 CN 1,2-Benzenedicarboxamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-N'-(1-phenylethyl)- (9CI) (CA INDEX NAME)



RN 328082-49-1 CAPLUS
 CN Cyclopentaneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 328082-50-4 CAPLUS
 CN Benzamide, 4-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-nitro- (9CI) (CA INDEX NAME)




RN 328082-51-5 CAPLUS

CN Cyclopropanecarboxamide, 2,2-dichloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-1-methyl- (9CI) (CA INDEX NAME)



RN 328082-52-6 CAPLUS


CN 1-Piperazinecarboxylic acid, 4-[5-[[[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]amino]carbonyl]-2-methoxyphenyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 328082-53-7 CAPLUS

CN 3-Pyrrolidinecarboxamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-oxo-1-(2-thienylmethyl)- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




RN 328082-55-9 CAPLUS

CN Benzamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-fluoro-(9CI) (CA INDEX NAME)




RN 328082-56-0 CAPLUS

CN Benzamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methyl-(9CI) (CA INDEX NAME)




RN 328082-57-1 CAPLUS

CN Benzamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-methyl-(9CI) (CA INDEX NAME)




RN 328082-58-2 CAPLUS


CN Benzamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(hydroxymethyl)-(9CI) (CA INDEX NAME)




RN 328083-64-3 CAPLUS
 CN 2H-Isoindole-2-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-1,3-dihydro-1,3-dioxo- (9CI) (CA INDEX NAME)




RN 328083-73-4 CAPLUS
 CN 4-Piperidinamine, 1-[(3,4-dichlorophenyl)methyl]-N-[(4-methylsulfonyl)phenyl]methyl- (9CI) (CA INDEX NAME)




RN 328083-75-6 CAPLUS
 CN 2-Thiazoleacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-phenyl- (9CI) (CA INDEX NAME)




RN 328083-76-7 CAPLUS
 CN 4-Thiazoleacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-phenyl- (9CI) (CA INDEX NAME)




RN 389062-07-1 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI)
 (CA INDEX NAME)




RN 479554-54-6 CAPLUS
 CN [1,1'-Biphenyl]-4-acetamide, N-[1-[(3,4-dihydroxyphenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-59-1 CAPLUS
 CN Benzeneacetamide, 4-bromo-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)



RN 479554-60-4 CAPLUS
 CN Benzeneacetamide, 4-amino-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-61-5 CAPLUS

CN Benzeneacetamide, 2-bromo-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-
(9CI) (CA INDEX NAME)


RN 479554-62-6 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-methyl- (9CI) (CA INDEX NAME)




RN 479554-63-7 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methyl- (9CI) (CA INDEX NAME)




RN 479554-64-8 CAPLUS

CN Benzeneacetamide, 3-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-hydroxy- (9CI) (CA INDEX NAME)




RN 479554-65-9 CAPLUS


CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-nitro- (9CI) (CA INDEX NAME)




RN 479554-66-0 CAPLUS
 CN Benzeneacetamide, 2-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-67-1 CAPLUS
 CN Benzeneacetamide, 4-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-69-3 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-nitro- (9CI) (CA INDEX NAME)




RN 479554-71-7 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,4-dimethoxy- (9CI) (CA INDEX NAME)






RN 479554-72-8 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-fluoro-4-hydroxy- (9CI) (CA INDEX NAME)




RN 479554-73-9 CAPLUS

CN 1,3-Benzodioxole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-75-1 CAPLUS

CN [1,1'-Biphenyl]-4-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-76-2 CAPLUS

CN Benzeneacetamide, 3,4-dichloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-82-0 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-methyl- (9CI) (CA INDEX NAME)




RN 479554-83-1 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(trifluoromethoxy)- (9CI) (CA INDEX NAME)




RN 479554-84-2 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-methoxy- (9CI) (CA INDEX NAME)




RN 479554-85-3 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(dimethylamino)- (9CI) (CA INDEX NAME)




RN 479554-87-5 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,4,5-trimethoxy- (9CI) (CA INDEX NAME)



RN 479554-89-7 CAPLUS


CN Benzeneacetamide, 3-amino-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479554-90-0 CAPLUS

CN 1-Naphthaleneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



RN 479554-91-1 CAPLUS


CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-hydroxy-3-methoxy- (9CI) (CA INDEX NAME)



RN 479554-92-2 CAPLUS

CN Benzeneacetamide, 3-[[[6-bromo-1-(2-propenyl)-2-naphthalenyl]oxy]methyl]-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 1-B

 $\text{--} \text{Cl}$


RN 479554-93-3 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-[(4-nitrophenyl)methoxy]- (9CI) (CA INDEX NAME)




RN 479554-94-4 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-fluoro-4-methoxy- (9CI) (CA INDEX NAME)




RN 479554-96-6 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-hydroxy- (9CI) (CA INDEX NAME)




RN 479554-97-7 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(phenylmethoxy)- (9CI) (CA INDEX NAME)




RN 479554-98-8 CAPLUS

CN [1,1'-Biphenyl]-4-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3'-nitro- (9CI) (CA INDEX NAME)




RN 479554-99-9 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,5-dimethyl- (9CI) (CA INDEX NAME)




RN 479555-00-5 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-iodo- (9CI) (CA INDEX NAME)




RN 479555-01-6 CAPLUS

CN Benzeneacetamide, 3-bromo-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-02-7 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methyl-3-nitro- (9CI) (CA INDEX NAME)




RN 479555-03-8 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-hydroxy-4-methoxy- (9CI) (CA INDEX NAME)




RN 479555-04-9 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-fluoro- (9CI) (CA INDEX NAME)




RN 479555-05-0 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-fluoro- (9CI) (CA INDEX NAME)




RN 479555-06-1 CAPLUS


CN Benzeneacetamide, 3-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-07-2 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-dimethyl- (9CI) (CA INDEX NAME)




RN 479555-09-4 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,4-difluoro- (9CI) (CA INDEX NAME)



RN 479555-10-7 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,4-difluoro- (9CI) (CA INDEX NAME)




RN 479555-11-8 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-difluoro- (9CI) (CA INDEX NAME)




RN 479555-12-9 CAPLUS

CN 3-Pyridineacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl] -
(9CI) (CA INDEX NAME)




RN 479555-13-0 CAPLUS

CN 2-Pyridineacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl] -
(9CI) (CA INDEX NAME)




RN 479555-14-1 CAPLUS

CN 3-Pyridineacetamide, 5-bromo-N-[1-[(3,4-dichlorophenyl)methyl]-4-
piperidinyl] - (9CI) (CA INDEX NAME)




RN 479555-15-2 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,4-
dimethoxy- (9CI) (CA INDEX NAME)



RN 479555-16-3 CAPLUS


CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-(phenylmethoxy)- (9CI) (CA INDEX NAME)




RN 479555-17-4 CAPLUS

CN 1-Naphthaleneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methyl- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




RN 479555-18-5 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-ethoxy- (9CI) (CA INDEX NAME)




RN 479555-19-6 CAPLUS

CN Benzeneacetamide, 4-butoxy-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-20-9 CAPLUS


CN 1H-Indole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)



RN 479555-21-0 CAPLUS


CN 2-Thiopheneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)






RN 479555-22-1 CAPLUS

CN Benzeneacetamide, 2,4-dichloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-23-2 CAPLUS

CN Benzeneacetamide, 2,6-dichloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-24-3 CAPLUS

CN 4-Thiazoleacetamide, 2-[(chloroacetyl)amino]-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-25-4 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,5-dimethoxy- (9CI) (CA INDEX NAME)




RN 479555-26-5 CAPLUS

CN 1H-Indole-3-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methoxy- (9CI) (CA INDEX NAME)




RN 479555-27-6 CAPLUS

CN 1H-Indole-3-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methoxy- (9CI) (CA INDEX NAME)




RN 479555-28-7 CAPLUS

CN 2-Naphthaleneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methoxy- (9CI) (CA INDEX NAME)




RN 479555-29-8 CAPLUS


CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(methylsulfonyl)- (9CI) (CA INDEX NAME)




RN 479555-30-1 CAPLUS
 CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,4,6-trimethyl- (9CI) (CA INDEX NAME)




RN 479555-31-2 CAPLUS
 CN 1H-Indole-3-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2-methyl-5-(1-methylethyl)- (9CI) (CA INDEX NAME)



RN 479555-32-3 CAPLUS
 CN 2H-Tetrazole-2-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-(1-pyrrolidinyl)- (9CI) (CA INDEX NAME)




RN 479555-33-4 CAPLUS
 CN 2H-Tetrazole-2-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-(4-methylphenyl)- (9CI) (CA INDEX NAME)




RN 479555-34-5 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-methoxy- (9CI) (CA INDEX NAME)




RN 479555-35-6 CAPLUS

CN Benzo[b]thiophene-3-acetamide, 5-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-36-7 CAPLUS

CN 4-Thiazoleacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methyl-2-phenyl- (9CI) (CA INDEX NAME)




RN 479555-37-8 CAPLUS

CN Benzo[b]thiophene-2-acetamide, 5-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-methyl- (9CI) (CA INDEX NAME)




RN 479555-38-9 CAPLUS

CN Benzo[b]thiophene-2-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-methyl- (9CI) (CA INDEX NAME)




RN 479555-39-0 CAPLUS

CN 1H-1,2,4-Triazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3-nitro- (9CI) (CA INDEX NAME)




RN 479555-40-3 CAPLUS

CN 1H-Pyrazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-5-methyl-3,4-dinitro- (9CI) (CA INDEX NAME)




RN 479555-41-4 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-(3-methylbutoxy)- (9CI) (CA INDEX NAME)




RN 479555-42-5 CAPLUS

CN 1H-Indole-5-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,3-dimethyl- (9CI) (CA INDEX NAME)




RN 479555-43-6 CAPLUS

CN 1H-Pyrazole-1-acetamide, 4-chloro-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-dimethyl- (9CI) (CA INDEX NAME)



RN 479555-44-7 CAPLUS


CN 1H-Pyrazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-dimethyl-4-nitro- (9CI) (CA INDEX NAME)



RN 479555-45-8 CAPLUS

CN 1H-Imidazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-2,4-dinitro- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A



RN 479555-46-9 CAPLUS
CN 1H-Imidazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-nitro- (9CI) (CA INDEX NAME)

PAGE 1-A




PAGE 2-A




RN 479555-47-0 CAPLUS

CN 1H-Pyrazole-1-acetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-3,5-dimethyl- (9CI) (CA INDEX NAME)




RN 479555-48-1 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-4-hexyl- (9CI) (CA INDEX NAME)




RN 479555-49-2 CAPLUS

CN Benzeneacetamide, 2-cyano-N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-65-2 CAPLUS

CN Benzeneacetamide, N-[1-[(2,5-dichlorophenyl)methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)




RN 479555-66-3 CAPLUS

CN Benzeneacetamide, N-[1-[(2,3-dichlorophenyl)methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)




RN 479555-67-4 CAPLUS


CN Benzeneacetamide, 4-fluoro-N-[1-[(4-fluorophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-68-5 CAPLUS
 CN Benzoic acid, 2-bromo-5-[[4-[(4-fluorophenyl)acetyl]amino]-1-piperidinyl]methyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 479555-69-6 CAPLUS
 CN Benzeneacetamide, 4-fluoro-N-[1-[(4-nitrophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)



RN 479555-70-9 CAPLUS
 CN Benzeneacetamide, N-[(1-[(3-benzoylphenyl)methyl]-4-piperidinyl)-4-fluoro- (9CI) (CA INDEX NAME)




RN 479555-74-3 CAPLUS
 CN Benzeneacetamide, 4-fluoro-N-[(1-[(4-methyl-3-nitrophenyl)methyl]-4-piperidinyl)-4- (9CI) (CA INDEX NAME)




RN 479555-75-4 CAPLUS

CN Benzeneacetamide, N-[1-[(3,4-dimethylphenyl)methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)




RN 479555-76-5 CAPLUS

CN Benzeneacetamide, 4-fluoro-N-[1-[(4-methoxy-3-methylphenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 479555-77-6 CAPLUS


CN [1,1'-Biphenyl]-2-carboxamide, 4'--[[4-[[4-(4-fluorophenyl)acetyl]amino]-1-piperidinyl]methyl]- (9CI) (CA INDEX NAME)



RN 479555-78-7 CAPLUS

CN Benzeneacetamide, N-[1-[[4-[[2,6-dichlorophenyl]methyl]sulfonyl]phenyl]methyl]-4-piperidinyl]-4-fluoro- (9CI) (CA INDEX NAME)






IT 68844-77-9, Astemizole

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (combination therapy component; prepn. and pharmaceutical combinations
 of [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3
 antagonists for treatment of asthma, allergic disease, or inflammation)

RN 68844-77-9 CAPLUS

CN 1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




IT 328082-07-1, N-[1-(3,4-Dichlorobenzyl)-4-piperidinyl]-N-[(6-methyl-2-pyridinyl)methyl]amine



RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (prepn. and pharmaceutical combinations of
 [(hetero)arylalkyl]piperidinyl amine, amide, or carbamate CCR3
 antagonists for treatment of asthma, allergic disease, or inflammation)

RN 328082-07-1 CAPLUS

CN 2-Pyridinemethanamine, N-[1-[(3,4-dichlorophenyl)methyl]-4-piperidinyl]-6-methyl- (9CI) (CA INDEX NAME)



GI



AB Title compds. I [wherein Z = CR₄R₅, CO, or CR₄R₅Z₁; Z₁ = alkylene, alkenylene, or CONH; R₁ = (un)substituted alkyl, alkenyl, (hetero)cycloalkyl, or (hetero)aryl; Q = O, S, NR₉, CO, CONR₉, NR₉CO, or CH=CH; m = 0-1; n = 0-6 with the proviso that when n = 0; then m = 0; R₂ and R₃ = independently H or alkyl; or CR₂R₃ = (alkyl)cycloalkyl; T = NR₁₀, CONR₁₀, NR₁₁CONR₁₀, or CONR₁₀R₁₁; X₁-X₄ = independently CH₂CHR₁₂ or CO; R₄ and R₅ = independently H or alkyl; R₆ = (un)substituted (hetero)aryl; R₉-R₁₁ = independently H, alkyl, haloalkyl, hydroxyalkyl, cycloalkyl(alkyl), or phenylalkyl; R₁₂ = independently (cyclo)alkyl or CO; or R₁₂ groups of X₁ and X₃ or X₄, or X₂ and X₃ or X₄ join to form CH₂CH₂, CH₂CH₂CH₂, CH₂OCH₂, or CH₂SCH₂; or pharmaceutically acceptable salts or solvates thereof] were prep'd. as cysteine-cysteine chemokine receptor 3 (CCR3) antagonists for use in pharmaceutical combinations with a histamine antagonist, steroid, leukotriene modulator, human cytokine, .beta.-agonist, phosphodiesterase inhibitor, or antibody (no data). For example, 1-(3,4-dichlorobenzyl)-4-piperidinamine.bul.2CF₃CO₂H was condensed with 2-(4-fluorophenyl)acetic acid to give N-[1-(3,4-dichlorobenzyl)-4-piperidinyl]-2-(4-fluorophenyl)acetamide (II). I are useful in combination therapy for the treatment of asthma, rhinitis, and other allergic or inflammatory conditions (no data).

L4 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:906175 CAPLUS

DN 138:14074

TI Preparation of benzo[g]quinoxalines for use against infectious diseases

IN Pato, Janos; Keri, Gyoergy; Oerfi, Laszlo; Waczek, Frigyes; Horvath, Zoltan; Banhegyi, Peter; Szabadkai, Istvan; Marosfalvi, Jenoe; Hegymegi-barakonyi, Balint; Szekelyhidi, Zsolt; Greff, Zoltan; Choidas, Axel; Bacher, Gerald; Daub, Henrik; Obert, Sabine; Kurtenbach, Alexander; Habenberger, Peter

PA Axxima Pharmaceuticals Ag, Germany; et al.

SO PCT Int. Appl., 237 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002094796	A2	20021128	WO 2002-EP5573	20020521
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,				

PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 2001-112289 A 20010518

US 2001-292325PP 20010522

US 2001-298902PP 20010619

EP 2001-115508 A 20010627

OS MARPAT 138:14074


IT 476637-76-0P, N,N'-Bis(1-benzylpiperidin-4-yl)benzo[g]quinoxaline-
 2,3-diamine

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)


(drug candidate; prepn. of benzo[g]quinoxalines for use against
 infectious diseases)

RN 476637-76-0 CAPLUS

CN Benzo[g]quinoxaline-2,3-diamine, N,N'-bis[1-(phenylmethyl)-4-piperidinyl]-
 (9CI) (CA INDEX NAME)



GI




AB The present invention relates to benzo[g]quinoxaline derivs. (shown as I; e.g. 2,3-bis(2-thienyl)benzo[g]quinoxaline and benzo[g]quinoxalin-2-yl(3-bromophenyl)amine), processes for manufg. said benzo[g]quinoxaline derivs., the use of the benzo[g]quinoxaline derivs. as pharmaceutically active agents, esp. for the prophylaxis and/or treatment of infectious diseases and opportunistic infections, diabetes, cancer, inflammation, as well as compns. contg. at least one benzo[g]quinoxaline deriv. and/or pharmaceutically acceptable salt thereof. Further, the present invention is directed to methods for preventing and/or treating of infectious

diseases, diabetes, cancer, and inflammation using the inventive benzo[g]quinoxaline derivs. The inventive benzo[g]quinoxaline derivs. exert their antiproliferative effect on *M. bovis* BCG and *M. tuberculosis* Erdmann at concns. between <<1 .mu.M and 32 .mu.M. In contrast, growth of *E. coli* XI-1 blue was not affected by benzo[g]quinoxaline derivs. at concns. >10 .mu.M. The benzo[g]quinoxaline compds. are able to inhibit HI virus replication up to 63% after 6 days at a concn. of 1 .mu.M. 5,10-Dibromo-2-(thiophen-3-yl)-3-(thiophen-2-yl)benzo[g]quinoxaline is able to decrease the activity of the herpes viral target UL-97 by 75%. Results for inhibition of HCMV target RICK for 5 I, of influenza replication for 7 I, of hepatitis B virus for 5 I, of TNF.alpha. signaling for 11 I, of human cellular protein kinases (Akt, Abl, PDGFR, Src) for 7 I, of A549 and Jurkat cells for 18 I, of human cellular protein kinase Akt known as a target for diabetes for 4 I, and of human protein kinases SRPK1 and SRPK2 (indicative of hepatitis B virus replication inhibition) for 8 and 1 I, resp., are tabulated. Results for activation of the insulin receptor InsR by 3 I, effect of 2 I on viability of Huh-5-2 replicon cells by the Alamar Blue toxicity assay, effect of 2 I on autonomous replication of hepatitis C virus replicons in the Huh-5-2 cell line by luciferase reporter assay, are tabulated. In I: R1 and R2 = -(CH₂)p-NH-(CH₂)n-R9, -(CH₂)s-S-(CH₂)m-R10, -(CH₂)m-O-(CH₂)p-R11, -(CH₂)r-R3, -CH:CH-R11, -(CH₂)m-CH(OH)(CH₂)p-R11, -(CH₂)q-R11, -R9, R10, -R12, -R13, etc. R3, R4, R5, R6, R7, and R8 = -H, -F, -Cl, -Br, -I, -SO₃H, -SO₃NH₂, -(CH₂)s-COOR16, -(CH₂)p-COOR17, -OR16, -SR16, -NR16R17, -OOCR16, -OOCR17, -NH-CO-R16, -NH-CO-R17, -CO-NH-R16, -CO-NH-R17, -NO₂, -N₃, -CN, -OCN, -NCO, -SCN, -NCS, CO-R16, CO-R17, -COCN, -CONR16R17, -SOR16, -SO₂R16, -SO₂R17, -SO₃R16, -SO₃R17, OCF3. R9, R10, and R11 = -CN, NR16R17, -NHR16, NHR17, etc. R12, R13, R14, and R15 = R3, R4, R5, R6, R16, R17, CH(CO₂R16)(CO₂R17), CH(CN)(CO₂R16), CH(CN)C(O)NHAr (Ar = R₁₄- and R₁₅-substituted phenyl); R16 and R17 = -H, -CH₃, -C₂H₅, -Pr, -CHMe₂, -Bu, -C₅H₁₁, -C₆H₁₃, -cyclo-C₆H₁₁, -cyclo-C₅H₉, -cyclo-C₄H₇, -cyclo-C₃H₅, -(CH₂)r-CHMe₂, -CHMeEt, -CMe₃, -CH:CH₂, -CH₂-CH:CH₂, Ph, --CH₂Ph, -C₂H₄Ph, -CH(CN)₂, -CF₃, -CCl₃, -CBr₃, -C₂F₅, -(CH₂)r-OH, -CH₂F, -CH₂Cl, -CH₂Br, -CH₂I, -CHF₂, -CHCl₂, -CHBr₂, -(CH₂)r-SH, -C₆H₄-CH₃, -C₆H₃Me₂, pyridyl, 2-pyrimidinyl, etc. M = 0-6, n = 0-6, p = 0-6, q = 0-6, r = 1-6, s = 0-6. Also claimed are the corresponding N-oxides in position 1 and/or 4 of these compds., the corresponding reduced forms of these compds. wherein the double bond in position 1 and/or 3 is hydrogenated, and pharmaceutically acceptable salts of I. About 42 example preps. and 406 compds. with characterization data are included. 1H-benzo[g]quinoxaline-2-one was prep'd. in 90% yield by dissolving 20 mmol 2,3-diaminonaphthalene in a mixt. of 5 mL DMF and 50 mL EtOH and adding 5 mL aq. soln. (50%) of glyoxalic acid and the mixt. was stirred for 2 h at reflux temp. The reaction mixt. was cooled to room temp. and the product was filtered, washed two times with Et₂O and dried.



L4 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:637660 CAPLUS
 DN 137:185501
 TI Preparation of quinazolines as specific inhibitors of type-13 matrix metalloprotease
 IN Andrianjara, Charles; Chantel-Barvian, Nicole; Gaudilliere, Bernard; Jacobelli, Henri; Ortwine, Daniel Fred; Patt, William Chester; Pham, Ly; Kostlan, Catherine Rose; Wilson, Michael William
 PA Warner-Lambert Company, USA
 SO PCT Int. Appl., 264 pp.
 CODEN: PIXXD2
 DT Patent

LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE	
PI	WO 2002064572	A1	20020822	WO 2002-EP1979	20020211	
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2001-268661PP	20010214
	US 2002193377	A1	20021219	US 2002-75954	20020213	
				US 2001-268661PP	20010214	
OS	MARPAT 137:185501					
IT	449209-88-5P					
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)					
	(MMP13 inhibitor; prepn. of quinazolines as specific inhibitors of type-13 matrix metalloprotease)					
RN	449209-88-5 CAPLUS					
CN	6-Quinazolinecarboxamide, 1,2,3,4-tetrahydro-1-methyl-2,4-dioxo-3-(phenylmethyl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)					



GI



AB Title compds. I [R1 = H, amino, alkyl, alkenyl, alkynyl, alkylamino, aryl, heterocycle, etc.; W = O, S, =N-R'; R' = alkyl, OH, CN; X1-3 = N, C-R6; R6 = H, alkyl, amino, alkylamino, etc.; Y = O, S, NH, N-alkyl; Z = O, S, NR7; R7 = H, alkyl, aryl, aryl, heteroaryl, etc.; n = 1-8; Z1 = alkyl; A = (non)arom., 5- or 6-membered monocycle comprising from 0 to 4 heteroatoms selected from N, O, S, etc.; m = 0-7; R2 = alkyl, halo, CN, NO2, SCF3, CF3, OCF3, etc.; R3 = H, alkyl, alkenyl, alkynyl, etc.] were prepd. Over 200 synthetic examples were provided. For instance, di-Me 4-aminoisophthalate was reacted with benzylisocyanate and heated to 95-100.degree. overnight to give Me 3-benzyl-2,4-dioxo-1,2,3,4-tetrahydroquinazoline-6-carboxylate which was saponified (dioxaneaq, LiOH, reflux) to give the carboxylic acid. This intermediate was coupled with benzylamine to afford II. Selected examples of I had IC50 = 2.25 - 0.001 .mu.M for MMP13 and IC50 > 100 .mu.M for MMP1, MMP2, MMP3, MMP7, MMP9, MMP12 and MMP14; II had IC50 = 0.193 .mu.M for MMP13. Compds. I are useful for the treatment of osteoarthritis and rheumatoid arthritis.

RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:123617 CAPLUS
 DN 136:183819
 TI Preparation of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors
 IN Wang, Wei-Bo; Curtin, Michael L.; Fakhoury, Stephen A.; Gwaltney, Stephen L.; Hasvold, Lisa A.; Hutchins, Charles W.; Li, Qun; Lin, Nan-Horng; Nelson, Lissa Taka Jennings; O'Connor, Steve; Sham, Hing L.; Sullivan, Gerard M.; Wang, Gary T.; Wang, Xilu
 PA USA
 SO U.S. Pat. Appl. Publ., 189 pp.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002019527	A1	20020214	US 2001-842391	20010425
				US 2000-200165PP	20000427

OS MARPAT 136:183819


IT 371761-79-4P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors)


RN 371761-79-4 CAPLUS

CN Benzonitrile, 4-[(1-methyl-1H-imidazol-5-yl)[(1-(phenylmethyl)-4-piperidinyl]amino)methyl]-2-(1-naphthalenyl)-, trihydrochloride (9CI) (CA INDEX NAME)



● 3 HCl

GI



II

AB Title compds. (I) were prep'd. Thus, 2-MeC₆H₄C₆H₃(CN)(CHO)-2,5 was condensed with 1-methyl-2-triethylsilyl-1H-imidazole (prepn. each given) and the product O-arylated to give title compd. II. Data for biol. activity of I were given.

L4 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:107318 CAPLUS

DN 136:151163

TI Preparation of indazole derivatives as JNK enzyme inhibitors

IN Bhagwat, Shripad S.; Satoh, Yoshitaka; Sakata, Steven T.

PA Signal Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 412 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002010137	A2	20020207	WO 2001-US23890	20010730
	WO 2002010137	C2	20030206		
		W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		
		RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG		US 2000-221799PP 20000731
	US 2002103229	A1	20020801	US 2001-910950	20010723
				US 2000-221799PP	20000731
EP	1313711	A2	20030528	EP 2001-957332	20010730
		R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		US 2000-221799PP 20000731
				WO 2001-US23890W	20010730

OS MARPAT 136:151163


IT 395107-63-8P, N-[1-Benzyl-4-piperidyl]-3-[5-(1H-1,2,4-triazol-3-yl)-1H-indazol-3-yl]benzamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of indazole derivs. as JNK enzyme inhibitors)

RN 395107-63-8 CAPLUS

CN Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]-3-[5-(1H-1,2,4-triazol-3-yl)-1H-indazol-3-yl]- (9CI) (CA INDEX NAME)




AB Indazole derivs., 3-R1A-5-R2-1H-indazoles (1), having activity as selective inhibitors of JNK are disclosed. In 1: A is a direct bond, -(CH₂)_a, -(CH₂)bCH:CH(CH₂)c-, or -(CH₂)bC.tplbond.C(CH₂)c-; R₁ is aryl, heteroaryl or heterocycle fused to Ph, each being optionally substituted with 1-4 R₃; R₂ is -R₃, -R₄, -(CH₂)bC(O)R₅, -(CH₂)bC(:O)OR₅, -(CH₂)bC(O)NR₅R₆, -(CH₂)bC(O)NR₅(CH₂)cC(O)R₆, -(CH₂)bNR₅C(O)R₆, -(CH₂)bNR₅C(O)NR₆R₇, -(CH₂)bNR₅R₆, -(CH₂)bOR₅, -(CH₂)bSODR₅ or -(CH₂)bSO₂NR₅R₆. A is 1-6; b and c are the same or different and are 0-4; d is 0-2. R₃ is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, substituted aryl, arylalkyl,

substituted arylalkyl, heterocycle, substituted heterocycle, heterocyclealkyl, substituted heterocyclealkyl, -C(O)OR8, -C(O)R8, -C(O)NR8R9, -C(O)NR8OR9, -SO2NR8R9, -NR8SO2R9, -CN, -NO2, -NR8R9, -NR8C(O)R9, -NR8C(O)(CH2)bOR9, -NR8C(O)(CH2)bR9, -O(CH2)bNR5R9, or heterocycle fused to Ph. R4 is alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, each being optionally substituted with 1-4 R3, or R4 is halogen or hydroxy. R5, R6 and R7 are the same or different and are H, alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, wherein each of R5, R6 and R7 are optionally substituted with 1-4 R3. R8 and R9 are the same or different and at each occurrence independently H, alkyl, aryl, arylalkyl, heterocycle, or heterocyclealkyl, or R8 and R9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R8, R9, and R8 and R9 taken together to form a heterocycle are optionally substituted with 1-4 R3 with the proviso that: when A is a direct bond and R1 is Ph, R2 is not Me, methoxy, C(O)CH3 or C(O)H; when A is a direct bond and R1 is 4-Me-Ph, R2 is not Me; when A is a direct bond and R1 is 4-F-Ph, R2 is not trifluoromethyl; when A is a direct bond or -C.tplbond.C- and R1 is Ph, R2 is not -COOEt; and when A is a direct bond and R1 is 6,7-dimethoxyisoquinolin-1-yl, R2 is not hydroxy. Such compds. have utility in the treatment of a wide range of conditions that are responsive to JNK inhibition. Thus, methods of treating such conditions are also disclosed, as are pharmaceutical compns. contg. one or more compds. of the above compds. Many of the claimed compds. have IC50 values $\text{IC50} \leq 0.5 \mu\text{M}$ in the JNK2 assay, e.g. 5-[3-(4-fluorophenyl)-1H-indazol-5-yl]-2H-1,2,3,4-tetrazole. Although the methods of prepn. are not claimed, >400 example prepn. are included.


L4	ANSWER 8 OF 13	CAPLUS	COPYRIGHT 2003 ACS on STN
AN	2002:72037	CAPLUS	
DN	136:134667		
TI	Preparation of mercaptopyrrolidinecarboxamides related compounds as inhibitors of endothelin-converting enzyme		
IN	Aebi, Johannes; Blum, Denise; Bur, Daniel; Chucholowski, Alexander; Dehmlow, Henrietta; Kitas, Eric Argirios; Loeffler, Bernd Michael; Obst, Ulrike; Wallbaum, Sabine		
PA	F. Hoffmann-La Roche A.-G., Switz.		
SO	PCT Int. Appl., 160 pp.		
	CODEN: PIXXD2		
DT	Patent		
LA	English		
FAN.CNT 1			
	PATENT NO.	KIND	DATE
	-----	-----	-----
PI	WO 2002006222	A1	20020124
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		WO 2001-EP7950 20010710
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG		EP 2000-114947 A 20000719
EP	1303485	A1	20030423
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		EP 2001-949485 20010710
			EP 2000-114947 A 20000719

BR 2001012580	A	20030617	WO 2001-EP7950 W 20010710 BR 2001-12580 20010710 EP 2000-114947 A 20000719 WO 2001-EP7950 W 20010710 US 2001-907135 20010717
US 2002049243	A1	20020425	
US 6541638	B2	20030401	EP 2000-114947 A 20000719
OS MARPAT 136:134667 IT 393158-11-7P	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (prepn. of mercaptopyrrolidinecarboxamides as inhibitors of endothelin-converting enzyme)		
RN 393158-11-7 CAPLUS			
CN 1-Pyrrolidinecarboxylic acid, 4-mercaptop-2-[[[1-(phenylmethyl)-4-piperidinyl]amino]carbonyl]-, phenylmethyl ester, (2S,4R)- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.



GI




AB Title compds. [I; R1 = H, alkylcarbonyl, arylcarbonyl; R2 = alkyl, alkylcycloalkyl, cycloalkyl, haloalkyl, carboxyalkyl, aryl, alkynyl, aryloxalkyl, heterocyclyl, etc.; A = COR3, CH(OH)R4, CONR5R6; R3, R4 = alkyl, aryl, arylalkynyl, aralkyl, arylalkenyl; R5 = H, alkyl, cycloalkyl, cycloalkylalkyl, carboxyalkyl, aralkyl; R6 = alkyl, alkylcarbonylalkyl, cyanoalkyl, hydroxyalkyl, aminocarbonylalkyl, aryl, etc.; m = 0-2; X = SO2, CO, CO2, SO2NH, CONR13; R13 = H, alkyl, aryl, carboxyalkyl], and dimers thereof, were prep'd. Thus, (2S,4R)-[[4-(4-methoxybenzylsulfanyl)-1-(naphthalene-2-sulfonyl)pyrrolidine-2-carbonyl]methylamino]acetic acid (prepn. given) in CH₂Cl₂ were treated with NMM, HOBT in CH₂Cl₂, EDCI in CH₂Cl₂, and o-toluidine in CH₂Cl₂; the soln. was shaken overnight to give a residue which was treated with Et₃SiH in CF₃CO₂H at 80.degree. for 1 h to give (2S,4R)-4-mercpto-1-(naphthalene-2-sulfonyl)pyrrolidine-2-carboxylic acid methyl(o-tolylcarbamoylmethyl)amide. I inhibited


endothelin converting enzyme with IC₅₀ = 5-1000 nM.
 RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2001:904170 CAPLUS
 DN 136:37519
 TI Synthesis and use of triazaspirodecanone derivatives as neurokinin receptor antagonists
 IN Galley, Guido; Godel, Thierry; Goergler, Annick; Hoffmann, Torsten; Kolczewski, Sabine; Roever, Stephan
 PA F. Hoffmann-La Roche AG, Switz.
 SO PCT Int. Appl., 90 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001094346	A1	20011213	WO 2001-EP6305	20010601
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CO, CU, CZ, DE, DK, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			EP 2000-112285 A	20000608
US	2002006932	A1	20020117	US 2001-861795	20010521
US	6482829	B2	20021119		
EP	1292596	A1	20030319	EP 2001-945242	20010601
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			EP 2000-112285 A	20000608
				WO 2001-EP6305 W	20010601
BR	2001011538	A	20030701	BR 2001-11538	20010601
				EP 2000-112285 A	20000608
				WO 2001-EP6305 W	20010601
OS	MARPAT 136:37519				
IT	968-86-5P 972-17-8P 380203-34-9P				
	RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)				
	(intermediate; synthesis and use of triazaspirodecanone derivs. as neurokinin receptor antagonists)				
RN	968-86-5 CAPLUS				
CN	4-Piperidinecarbonitrile, 4-(phenylamino)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)				



RN 972-17-8 CAPLUS
 CN 4-Piperidinecarbonitrile, 4-[(2-methylphenyl)amino]-1-(phenylmethyl)-
 (9CI) (CA INDEX NAME)



RN 380203-34-9 CAPLUS
 CN 4-Piperidinemethanamine, 4-[(2-methylphenyl)amino]-1-(phenylmethyl)- (9CI)
 (CA INDEX NAME)



GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. I [R1 = H, alkyl, alkenyl, Ph, $(CH_2)_m$ -non arom. heterocyclyl, $(CH_2)_m$ -heteroaryl, $(CH_2)_m$ -carboxamide, $(CH_2)_m-C(O)alkyl$, etc.; R2 = H, alkyl, halo, alkoxy; R3 = alkyl, alkoxy, halo, CF3; X = N-, C:, CH; X1/X2 = H, OH, alkoxy or may be together an oxo group; Y1/Y2 = H, alkyl, $(CH_2)_m$ -Ph or may be together an oxo group; Z = bond, CH2, C(O); m = 0 - 4; n = 2 - 3; p = 0 - 2] were prep'd. Over 160 synthetic examples were disclosed. For example, 8-(3,5-bistrifluoromethylbenzoyl)-1-phenyl-1,3,8-triazaspiro[4.5]decan-4-one was reacted with 2-chloro-4,6-dimethoxy-1,3,5-


triazine (1,2-dimethoxyethane, NaH, 100.degree.C, 1 h) to give II. II had pKi = 8.66 for the NK-1 receptor. I are useful in the treatment of diseases related to NK-1 receptor antagonists.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2001:827020 CAPLUS
 DN 136:294764
 TI Synthesis of 2-(2,3-dimethoxyphenyl)-4-(aminomethyl)imidazole analogues and their binding affinities for dopamine D2 and D3 receptors
 AU Huang, Yunsheng; Luedtke, Robert R.; Freeman, Rebekah A.; Wu, Li; Mach, Robert H.
 CS Department of Radiology-PET Center, Wake Forest University School of Medicine, Winston-Salem, NC, 27157, USA
 SO Bioorganic & Medicinal Chemistry (2001), 9(12), 3113-3122
 CODEN: BMECEP; ISSN: 0968-0896
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 OS CASREACT 136:294764
 IT 407610-25-7P 407610-26-8P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (prepn. and dopamine D2 and D3 receptor affinity of
 2-(2,3-dimethoxyphenyl)-1H-imidazole-4-methanamine derivs.)
 RN 407610-25-7 CAPLUS
 CN 4-Piperidinamine, N-[(2-(2,3-dimethoxyphenyl)-1H-imidazol-4-yl)methyl]-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 407610-26-8 CAPLUS
 CN 4-Piperidinamine, N-[(2-(5-bromo-2,3-dimethoxyphenyl)-1H-imidazol-4-yl)methyl]-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



AB A series of 2-(2,3-dimethoxyphenyl)-4-(aminomethyl)imidazole derivs. was prep'd. and their affinity for dopamine D2 and D3 receptors was measured using in vitro binding assays. Several oxadiazole analogs were also prep'd. and tested for their affinity for dopamine D2 and D3 receptors. The results of receptor binding studies indicated that the

incorporation of an imidazole moiety between the Ph ring and the basic nitrogen did not significantly increase the selectivity for dopamine D3 receptors, whereas the incorporation of an oxadiazole at the same region resulted in a total loss of affinity for both dopamine receptor subtype binding sites. The most selective compd. in this series is 6,7-dimethoxy-2-[[2-(2,3-dimethoxyphenyl)-1H-imidazol-4-yl]methyl]-1,2,3,4-tetrahydroisoquinoline, which has a D3 receptor affinity of 21 nM and a 7-fold selectivity for D3 vs. D2 receptors. The binding affinity for .sigma.1 and .sigma.2 receptors was also measured, and the results showed that several analogs were selective .sigma.1 receptor ligands.

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4	ANSWER 11 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN			
AN	2000:260283 CAPLUS			
DN	132:293757			
TI	Preparation of novel 4,5-dihydroisoxazole derivatives and their use as pharmaceuticals for T cell-mediated diseases			
IN	Freyne, Eddy Jean Edgard; Andres-Gil, Jose Ignacio; Deroose, Frederik Dirk; Petit, Davy Petrus Franciscus Maria; Matesanz-Ballesteros, Maria Encarnacion; Alvarez Escobar, Rosa Maria			
PA	Janssen Pharmaceutica N.V., Belg.			
SO	PCT Int. Appl., 108 pp. CODEN: PIXXD2			
DT	Patent			
LA	English			
FAN.CNT 1				
	PATENT NO.	KIND	DATE	APPLICATION NO. DATE
-----	-----	-----	-----	-----
PI	WO 2000021959	A1	20000420	WO 1999-EP7803 19991007
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			EP 1998-203394 A 19981009
	CA 2346396	AA	20000420	CA 1999-2346396 19991007
				EP 1998-203394 A 19981009
				WO 1999-EP7803 W 19991007
	EP 1119568	A1	20010801	EP 1999-953847 19991007
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
				EP 1998-203394 A 19981009
				WO 1999-EP7803 W 19991007
	JP 2002527438	T2	20020827	JP 2000-575865 19991007
				EP 1998-203394 A 19981009
				WO 1999-EP7803 W 19991007
	AU 763460	B2	20030724	AU 2000-10393 19991007
				EP 1998-203394 A 19981009
				WO 1999-EP7803 W 19991007
	US 6583141	B1	20030624	US 2001-807149 20010406
				EP 1998-203394 A 19981009
				WO 1999-EP7803 W 19991007
OS	MARPAT 132:293757			


IT 264604-07-1P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (target compd.; prepn. of dihydroisoxazole derivs. as antiproliferatives and immunomodulators)


RN 264604-07-1 CAPLUS

CN 5-Isoxazolecarboxamide, 4,5-dihydro-N-[4-[phenyl[1-(phenylmethyl)-4-piperidinyl]amino]methyl]phenyl]-3-(3-pyridinyl)- (9CI) (CA INDEX NAME)



PAGE 1-A



PAGE 2-A



GI



AB The invention concerns title compds. I and their N-oxides, pharmaceutically acceptable addn. salts, quaternary ammonium salts, and stereochem. isomeric forms [wherein m, n, p = 0 or 1; R1 = (un)substituted pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl or phenyl; B = amide, ketone, or **oxadiazole**; D = (un)substituted aryl or heterocyclyl; Q = bond, CO, (un)substituted NH, CONH, CH₂, CH(:CH₂), C(:NH), SO, SO₂, 3-oxobut enyl, pyrazole, isoxazole, or thiazole nucleus; L = (un)substituted aryl or heteroaryl; R2, R3 = H, halo, C₁-6 alkyloxy, or (un)substituted C₁-6 alkyl]. Also disclosed is a process for their prepn., compns. comprising them, and their medical use. The compds. show growth inhibitory activity against T cell blasts and keratinocytes in vitro. The compds. are claimed for use in the treatment of prevention of rheumatic, arthritic, and inflammatory diseases, psoriasis, T cell leukemia, transplant rejection, and graft-vs.-host disease. For instance, base-catalyzed cycloaddn. of N-hydroxy-3-pyridinecarboximidoyl chloride with Me 2-propenoate gave 98% Me 4,5-dihydro-3-(3-pyridinyl)-5-isoxazolecarboxylate, which was amidated with (4-aminophenyl)phenylmethanone to give 58% title compd. II. At a concn. of 10-6 M, II gave 81% inhibition of T cell blast formation in human whole blood.


RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1999:64782 CAPLUS
 DN 130:139366
 TI Preparation of 6-azauracil derivatives as IL-5 biosynthesis inhibitors
 IN Lacrampe, Jean Fernand Armand; Freyne, Eddy Jean Edgard; Venet, Marc
 Gaston; Boeckx, Gustaaf Maria
 PA Janssen Pharmaceutica N.V., Belg.
 SO PCT Int. Appl., 83 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9902505	A1	19990121	WO 1998-EP4191	19980707
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HR, HU, ID, IL, IS, JP, KE,				


KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,
 MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
 TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES,
 FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI,
 CM, GA, GN, ML, MR, NE, SN, TD, TG

AU 9889738	A1	19990208	EP 1997-202118 A 19970710
AU 742145	B2	20011220	AU 1998-89738 19980707
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
EP 1000040	A1	20000517	EP 1998-941299 19980707
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, SI, LT, LV, FI, RO			
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
EE 200000016	A	20001016	EE 2000-20000001619980707
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
NZ 502180	A	20001124	NZ 1998-502180 19980707
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
TW 496865	B	20020801	TW 1998-87111014 19980708
			EP 1997-202118 A 19970710
ZA 9806089	A	20000110	ZA 1998-6089 19980709
			EP 1997-202118 A 19970710
BR 9811678	A	20000919	BR 1998-11678 19980710
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
HR 2000000003	A1	20001231	HR 2000-3 20000105
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
NO 2000000063	A	20000310	NO 2000-63 20000106
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
US 2002072603	A1	20020613	US 2001-891888 20010626
			EP 1997-202118 A 19970710
			WO 1998-EP4191 W 19980707
			US 2000-462320 B120000105
OS	MARPAT 130:139366		
IT	219979-02-9P 219979-22-3P		
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)		
	(prepn. of 6-azauracil derivs. as IL-5 biosynthesis inhibitors)		
RN	219979-02-9 CAPLUS		
CN	5-Thiazolecarboxamide, 2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]methyl]-4-phenyl-N-[1-(phenylmethyl)-4-piperidinyl] - (9CI) (CA INDEX NAME)		




RN 219979-22-3 CAPLUS

CN 5-Thiazoleacetamide, 2-[(4-chlorophenyl)[2,6-dichloro-4-(4,5-dihydro-3,5-dioxo-1,2,4-triazin-2(3H)-yl)phenyl]methyl]-4-phenyl-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB RZCR1(XR2)R3 [I; R= 3,5-dioxo-1,2,4-triazin-2(3H)-yl; R1 = H, halo, alkyl, alkoxy, etc.; R2 = CONH₂, (un)substituted alkyl, (hetero)aryl, etc.; R3 = (un)substituted Ph; X = bond, O, s, (alkyl)imino; Z = (un)substituted phenylene] were prep'd. Thus, title compd. II (R4 = Cl) was etherified by Me₂CHCH₂OH to give II (R4 = OCH₂CHMe₂). Data for biol. activity of I were given.

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN

AN 1994:30773 CAPLUS

DN 120:30773

TI Oxadiazole derivatives having acetylcholinesterase-inhibitory and muscarinic receptor agonist activity

IN Takasugi, Hisashi; Kuno, Atsushi; Ohkubo, Mitsuru

PA Fujisawa Pharmaceutical Co., Ltd., Japan

SO PCT Int. Appl., 149 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9313083	A1	19930708	WO 1992-JP1658	19921218
	W: AU, CA, HU, JP, KR, RU, US			GB 1991-27533	19911231
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			GB 1992-20904	19921005
AU	9331714	A1	19930728	AU 1993-31714	19921218
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
EP	619814	A1	19941019	EP 1993-900416	19921218
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE			GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
JP	07502529	T2	19950316	JP 1992-511547	19921218
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
US	5622976	A	19970422	US 1994-244904	19940624
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218

L6 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 1998:631744 CAPLUS
DN 129:310895
TI Benzamide compounds and their use as neovascularization inhibitors
IN Inaba, Takayuki; Tada, Hiroki; Iwamura, Hiroyuki

PA Japan Tobacco, Inc., Japan
 SO Jpn. Kokai Tokkyo Koho, 106 pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese
 FAN CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10259176	A2	19980929	JP 1997-84463	19970317
				JP 1997-84463	19970317


OS MARPAT 129:310895

IT 214846-51-2P



RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (Benzamide compds. and their use as neovascularization inhibitors)

RN 214846-51-2 CAPLUS

CN 2H-1,4-Benzoxazine-7-carboxamide, 3,4-dihydro-3-oxo-4-(phenylmethyl)-N-[1-(phenylmethyl)-4-piperidinyl] - (9CI) (CA INDEX NAME)



GI



AB The inhibitors contain benzamides I [R1 = H, NO₂, halo, cyano, lower alkoxy, NR₁₁R₁₂ (R₁₁, R₁₂ = H, acyl); R₂ = H, NO₂, halo, OR₁₃ (R₁₃ = lower alkyl, aralkyl, cycloalkyl); R₃ = X₃(CH₂)_mR₁₄ (R₁₄ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, (un)substituted lower alkyl, cycloalkyl, acyl, alkenyl, H; X₃ = O, NHCO, OSO₂, NR₁₇ (R₁₇ = H, lower alkyl); m = 0-5], II (R₁₅, R₁₆ = H, lower alkoxy, amino, lower alkyl, CO₂H, OH); R₂ and R₃ may be bonded to form a condensed 1,3-oxazole ring; R₄ = H, OR₁₉ (R₁₉ = lower alkyl, aralkyl, cycloalkyl); R₃ and R₄ may be bonded to form a condensed 1,3-oxazole, 1,4-oxazine, or pyrimidine ring; R₅ = H, NO₂, alkenyl, oxazole, NR₂₈ (R₂₈ = H, acyl, lower alkoxy carbonyl); R₆ = H, (un)substituted lower alkyl; R₅ and R₆ may be bonded to form a condensed pyrimidine, diazepine, or pyridine ring; R₇ = H, lower alkoxy; R₈ = X₄(CH₂)_tR₃₀ [X₄ = O, CH₂, CO, or CONH, OSO₂, SO₂NH, NR₃₁ (R₃₁ = H, lower alkyl, aralkyl), direct bond], t = 0-5; R₃₀ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, H, OH, halo, lower alkyl, lower alkoxy, cycloalkyl, acyl, cyano, neoplasms, etc. IC₅₀ of 4-benzyloxy-N-(4-benzyloxyphenyl)-3-methoxybenzamide (prepn. given) against bFGF- or VEGF-induced proliferation of HUVEC was 0.85 .mu.M.

L10 ANSWER 18 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
AN 1998:682229 CAPLUS

DN 129:302552
 TI Preparation of 1,4-disubstituted cyclic amine derivatives as serotonin antagonists
 IN Kitazawa, Noritaka; Ueno, Kohshi; Takahashi, Keiko; Kimura, Teiji; Sasaki, Atsushi; Kawano, Koki; Okabe, Tadashi; Komatsu, Makoto; Matsunaga, Manabu; Kubota, Atsuhiko
 PA Eisai Co., Ltd., Japan
 SO PCT Int. Appl., 635 pp.
 CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9843956	A1	19981008	WO 1998-JP1481	19980331
	W: AU, CA, CN, HU, JP, KR, MX, NO, NZ, RU, US			JP 1997-98433	A 19970331
	RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			JP 1997-366764	A 19971226
	AU 9865209	A1	19981022	AU 1998-65209	19980331
	AU 748038	B2	20020530	JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	ZA 9802707	A	19991020	WO 1998-JP1481	W 19980331
	EP 976732	A1	20000202	JP 1997-98433	A 19970331
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			EP 1998-911137	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	NZ 337651	A	20020426	WO 1998-JP1481	W 19980331
				NZ 1998-337651	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	RU 2203275	C2	20030427	WO 1998-JP1481	W 19980331
				RU 1999-123039	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	US 6448243	B1	20020910	WO 1998-JP1481	W 19980331
				US 1999-367227	19990811
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	NO 9904720	A	19991130	WO 1998-JP1481	W 19980331
				NO 1999-4720	19990928
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	US 2002086999	A1	20020704	WO 1998-JP1481	W 19980331
				US 2001-846259	20010502
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
				WO 1998-JP1481	W 19980331
	US 2002019531	A1	20020214	US 1999-367227	A319990811
	US 6579881	B2	20030617	US 2001-859517	20010518
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
				WO 1998-JP1481	W 19980331
				US 1999-367227	A319990811


OS MARPAT 129:302552

IT 202859-14-1P 214611-21-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (prepn. of 1,4-disubstituted cyclic amine derivs. as serotonin antagonists)


RN 202859-14-1 CAPLUS

CN 4-Piperidinamine, N-(3-methoxyphenyl)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 214611-21-9 CAPLUS

CN 4-Piperidinamine, N-(3-bromophenyl)-1-[2-(4-fluorophenyl)ethyl]- (9CI) (CA INDEX NAME)




IT 131587-28-5

RL: RCT (Reactant); RACT (Reactant or reagent)
 (prepn. of 1,4-disubstituted cyclic amine derivs. as serotonin antagonists)

RN 131587-28-5 CAPLUS

CN 4-Piperidinamine, N-(3-fluorophenyl)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



L4 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1994:30773 CAPLUS
 DN 120:30773
 TI Oxadiazole derivatives having acetylcholinesterase-inhibitory
 and muscarinic receptor agonist activity
 IN Takasugi, Hisashi; Kuno, Atsushi; Ohkubo, Mitsuru
 PA Fujisawa Pharmaceutical Co., Ltd., Japan
 SO PCT Int. Appl., 149 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9313083	A1	19930708	WO 1992-JP1658	19921218
	W: AU, CA, HU, JP, KR, RU, US			GB 1991-27533	19911231
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			GB 1992-20904	19921005
	AU 9331714	A1	19930728	AU 1993-31714	19921218
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
	EP 619814	A1	19941019	EP 1993-900416	19921218
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE			GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
	JP 07502529	T2	19950316	JP 1992-511547	19921218
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218
	US 5622976	A	19970422	US 1994-244904	19940624
				GB 1991-27533	19911231
				GB 1992-20904	19921005
				WO 1992-JP1658	19921218

Patel

<10/13/2003>


OS MARPAT 120:30773

IT 151097-86-8P 151097-87-9P 151307-60-7P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (prep. and acetylcholinesterase inhibitory and muscarinic receptor
 agonist activity of)

RN 151097-86-8 CAPPLUS


CN 1,2,4-Oxadiazole-3-carboxamide, 5-(1-azabicyclo[2.2.2]oct-3-yl)-N-[1-(phenylmethyl)-4-piperidinyl]-, dihydrochloride (9CI) (CA INDEX NAME)



● 2 HCl

RN 151097-87-9 CAPPLUS

CN 1,2,4-Oxadiazole-3-carboxamide, 5-(1-azabicyclo[2.2.2]oct-3-yl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 151307-60-7 CAPPLUS

CN 1,2,4-Oxadiazole-3-carboxamide, 5-(4-nitrophenyl)-N-[1-(phenylmethyl)-4-piperidinyl]-, (2E)-2-butenedioate (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 151307-59-4

CMF C21 H21 N5 O4




CM 2

10069215.2

CRN 110-17-8
 CMF C₄H₄O₄


Double bond geometry as shown.



AB The title compds. R1QZXAM [A = direct bond, lower alkylene, lower alkynylene; M = (un)substituted heterocyclic group contg. .gtoreq.1 N atom(s); Q = oxadiazolediyl; R1 = lower alkyl, (un)substituted heterocyclic group, (un)substituted aryl, (un)substituted arylalkyl, (un)substituted aralkenyl; X = direct bond, CONR4, R8CN; R4 = H, alkyl; R8 = HO, protected HO group, CO, NHCO; Z = direct bond, vinyl (sic)], useful for the treatment of central nervous system disorders (e.g., amnesia, Alzheimer's disease, vascular dementia, etc.) mode data, are prep'd. Thus, 3-ethoxycarbonyl-5-(quinuclidin-3-yl)-1,2,4-oxadiazole and 1-benzyl-4-(2-aminoethyl)piperidine were heated together in soln. at 100.degree. for 2 h and treated with an ethanolic soln. of HCl, producing 5-(quinuclidin-3-yl)-3-[(2-(1-benzylpiperidin-4-yl)ethyl)carbamoyl]-1,2,4-oxadiazole dihydrochloride, m.p. 210.degree. (decompn.).

CRN 110-17-8
CMF C4 H4 O4

Double bond geometry as shown.



AB The title compds. R1QZXAM [A = direct bond, lower alkylene, lower alkynylene; M = (un)substituted heterocyclic group contg. .gtoreq.1 N atom(s); Q = oxadiazolediyl; R1 = lower alkyl, (un)substituted heterocyclic group, (un)substituted aryl, (un)substituted arylalkyl, (un)substituted aralkenyl; X = direct bond, CONR4, R8CN; R4 = H, alkyl; R8 = HO, protected HO group, CO, NHCO; Z = direct bond, vinyl (sic)], useful for the treatment of central nervous system disorders (e.g., amnesia, Alzheimer's disease, vascular dementia, etc.) mode data, are prep'd. Thus, 3-ethoxycarbonyl-5-(quinuclidin-3-yl)-1,2,4-oxadiazole and 1-benzyl-4-(2-aminoethyl)piperidine were heated together in soln. at 100.degree. for 2 h and treated with an ethanolic soln. of HCl, producing 5-(quinuclidin-3-yl)-3-[[2-(1-benzylpiperidin-4-yl)ethyl]carbamoyl]-1,2,4-oxadiazole dihydrochloride, m.p. 210.degree. (decompn.).

=> d 16 fbib hitstr abs total

L6 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2003:319886 CAPLUS
DN 138:338155

TI Preparation of oxadiazolyl-biphenylcarboxamides as p38 kinase inhibitors
IN Angell, Richard Martyn; Bamborough, Paul; Cockerill, George Stuart; Smith, Kathryn Jane; Walker, Ann Louise

PA Glaxo Group Limited, UK
SO PCT Int. Appl., 44 pp.
CODEN: PIXXD2

DT Patent
LA English
FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003033482	A1	20030424	WO 2002-EP11574	20021016
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	GB 2001-24932 A 20011017

OS MARPAT 138:338155
IT 515143-78-9P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of oxadiazolyl-biphenylcarboxamides as p38 kinase inhibitors)

RN 515143-78-9 CAPLUS

CN [1,1'-Biphenyl]-4-carboxamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]-2'-methyl-5'-(5-methyl-1,3,4-oxadiazol-2-yl)- (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title compds. [I; X = a bond, (un)substituted Ph; R1 = (un)substituted 5-7 membered heterocyclyl, 5-7 membered heteroaryl, fused bicycyl; R2 = H, alkyl, (CH₂)pcycloalkyl; or when X = a bond and m and n are both zero, NR₁R₂ = 5-6 membered heterocyclyl optionally contg. one addnl. heteroatom selected from O and N which can be optionally substituted by alkyl; R3 = II (wherein R₄ = H, alkyl); U = Me, halo; V, Y = H, Me, halo; m, n = 0-2; m + n = 0-4; p = 0-1; r = 0-2; with the provisos], useful as pharmaceuticals, particularly as p38 kinase inhibitors, were prepd. E.g., a 6-step synthesis of the carboxamide III, starting from 3-bromo-4-methylbenzoic acid, was given.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2003:42265 CAPLUS

DN 138:106699

TI Preparation of (indazolyl)benzimidazoles and analogs as tyrosine and serine/threonine kinase inhibitors

IN Renhowe, Paul A.; Shafer, Cynthia M.; McBride, Chris; Silver, Joel; Pecchi, Sabina; Machajewski, Tim; McCrea, Bill; Poon, Daniel; Thomas, Teresa

PA Chiron Corporation, USA

SO PCT Int. Appl., 435 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003004488	A1	20030116	WO 2002-US20844	20020702
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,			

LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL,
 PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG,
 US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,
 PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
 NE, SN, TD, TG

US 2001-302791PP 20010703

OS MARPAT 138:106699


IT **485837-39-6P**, N-(1-Benzylpiperidin-4-yl)-2-(5-methoxy-1H-indazol-3-yl)-1H-benzimidazol-6-amine **485837-40-9P**, N-(1-Benzylpiperidin-4-yl)-2-(6-fluoro-1H-indazol-3-yl)-1H-benzimidazol-6-amine
485841-19-8P, N-(1-Benzylpiperidin-4-yl)-3-[6-(1,4'-bipiperidin-1'-yl)-1H-benzimidazol-2-yl]-1H-indazol-5-amine

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(kinase inhibitor; prepn. of (indazolyl)benzimidazole kinase inhibitors by cyclizing indazolyl aldehydes or ketones with phenylenediamines)


RN 485837-39-6 CAPLUS

CN 1H-Benzimidazol-5-amine, 2-(5-methoxy-1H-indazol-3-yl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 485837-40-9 CAPLUS

CN 1H-Benzimidazol-5-amine, 2-(6-fluoro-1H-indazol-3-yl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 485841-19-8 CAPLUS

CN 1H-Indazol-5-amine, 3-(5-[1,4'-bipiperidin]-1'-yl-1H-benzimidazol-2-yl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB Title compds. I [wherein Z1-Z4 = C independently C or N; R1 = H, F, Cl, or Br; R2 = H, F, Cl, Br, CN, NO₂, or (un)substituted CO₂H, NH₂, CONH₂, NHCONH₂, etc.; R3 = H, F, Cl, Br, or (un)substituted alkoxy; R4, R9, and R10 = H; R5 and R8 = independently H, F, Cl, or (un)substituted alkyl, alkoxy, NH₂, heterocyclyl, etc.; R6 and R7 = independently H, F, Cl, Br, CF₃, CO₂H, or (un)substituted alkyl, (heterocyclyl)alkoxy, arylalkoxy, alkoxyalkoxy, (heterocyclyl)heterocyclyl, arylheterocyclyl, heterocyclyoxy, aryloxy, NH₂, CONH₂, etc.; or R5 is absent if Z1 = N; or R6 is absent if Z2 = N; or R7 is absent if Z3 = N; or R8 is absent if Z4 = N; with the proviso that at least one of R1, R2, R3, R5, R6, R7, or R8 .noteq. H; and tautomers and pharmaceutically acceptable salts thereof] were prep'd. as tyrosine and serine/threonine kinase inhibitors. For example, dimerization of indazole-3-carboxylic acid with PO₃ followed by addn. of 1,2-phenylenediamine in toluene gave 3-(1H-benzimidazol-2-yl)-1H-indazole. Seven hundred twenty-eight exemplary compds. were assayed for serine/threonine kinase activity in vitro, and the majority displayed an IC₅₀ value of less than 10 .mu.M with respect to VEGFR1, Flk-1, bFGF, Tie-2, CHK-1, cdc2, GSK-3, NEK-2, and PDGF.

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2001:115087 CAPLUS
DN 134:163028
TI Solid phase synthesis of **oxazoles** and thiazoles

IN Bunin, Barry A.; Tushup, Steven P.
 PA ChemRx Advanced Technologies, Inc., USA
 SO PCT Int. Appl., 43 pp.
 CODEN: PIXXD2

DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001010798	A1	20010215	WO 2000-US21051	20000802
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			US 1999-147451PP 19990804


OS CASREACT 134:163028; MARPAT 134:163028

IT 325709-13-5P


RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)
 (solid phase synthesis of oxazoles and thiazoles)

RN 325709-13-5 CAPLUS

CN 5-Oxazolecarboxamide, 2-[4-(hexyloxy)phenyl]-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB Title compds. [I; X1 = O, S; R1 = R3, NR3R4, NR3C(:NR4)NR3R4, NR3CONR3R4; R3 = alkyl, (substituted) aryl(alkyl), polycycloaryl(alkyl), heatheroaryl(alkyl), etc.; R4 = H, alkyl; X2 = piperidinyl, pyrrolidinyl, CHR8NR7, C6H4CH2NR7; R7 = H, alkyl; R8 = H, (substituted) alkyl, cycloalkylalkyl, heterocycloalkylalkyl, aralkyl, heteroarylalkyl, polycycloaryl(alkyl), heteropolycycloarylalkyl; R2 = CH2OH, CONR9R10, CO2R11; R9-R11 = H, alkyl, cycloalkylalkyl, heterocyclalkyl, aryl(alkyl), heteroaryl(alkyl), polycycloaryl(alkyl), heteropolycyclo(alkyl), etc.], and arrays thereof were prep'd. by (1) treatment of 4-(SSCH2S)C6H4O2CX2H (SS = solid support; X2 as above) with I (R2X2 = OH) to give supported intermediates (II; variables as above), and (2) treatment of II or arrays thereof with reducing agents, amines, or alcs. Thus, II [SS = Merrifield resin; X2 = CH(CH2Ph)NH] was swelled in CH2Cl2 and added to a mixt. of 2-p-tolyloxazole-4-carboxylic acid and DIC in CH2Cl2 followed by stirring, addn. of dimethylaminopyridine, and stirring for 15 h to give functionalized resin. This was suspended in dioxane and stirred 10-24 h with BuNH2 to give 2-p-tolyl-4-carboxylic acid (1-butylcarbamoyl-2-phenylethyl)amide.

RE.CNT 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L6 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2000:227639 CAPLUS
 DN 132:251141
 TI Preparation of oxazole compounds as prostaglandin E2 agonists or antagonists
 IN Hattori, Kouji; Tanaka, Akira; Kono, Yutaka; Nakazato, Shoko
 PA Fujisawa Pharmaceutical Co., Ltd., Japan
 SO PCT Int. Appl., 121 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000018744	A1	20000406	WO 1999-JP5212	19990924
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			AU 1998-6176	A 19980925
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			AU 1999-9822	A 19990419
CA	2345474	AA	20000406	CA 1999-2345474	19990924
				AU 1998-6176	A 19980925
				AU 1999-9822	A 19990419
AU	9957590	A1	20000417	WO 1999-JP5212	W 19990924
				AU 1999-57590	19990924
				AU 1998-6176	A 19980925
				AU 1999-9822	A 19990419
BR	9914451	A	20010522	WO 1999-JP5212	W 19990924
				BR 1999-14451	19990924
				AU 1998-6176	A 19980925
				AU 1999-9822	A 19990419

EP 1115712	A1	20010718	WO 1999-JP5212 W 19990924
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			EP 1999-944806 19990924
JP 2002525361	T2	20020813	AU 1998-6176 A 19980925 AU 1999-9822 A 19990419 WO 1999-JP5212 W 19990924 JP 2000-572204 19990924
US 6437146	B1	20020820	AU 1998-6176 A 19980925 AU 1999-9822 A 19990419 WO 1999-JP5212 W 19990924 US 2001-787433 20010420
OS	MARPAT 132:251141		
IT	262594-98-9P 262595-76-6P		
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (prepn. of oxazole compds. as prostaglandin E2 agonists or antagonists)			
RN	262594-98-9 CAPLUS		
CN	Benzamide, 3-[(1S,2R)-2-(4,5-diphenyl-2-oxazolyl)cyclopentyl]methyl]-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)		



Absolute stereochemistry.



RN	262595-76-6 CAPLUS
CN	Benzamide, 3-[(2-(4,5-diphenyl-2-oxazolyl)-2-cyclohexen-1-yl)methyl]-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



II

AB Oxazole compds. of formula I [R1 = aryl which may be substituted with halogen(s); R2 = aryl which may be substituted with halogen(s), X = single bond, or SO₂, R3, R4 = H or suitable substituent, (wherein X is neither R3 nor R4 is hydrogen), R3 and R4 may be linked together to form an N-contg. heterocyclic group which may be substituted with one or more suitable substituent(s), R5 = H, etc., A1 = lower alkylene or single bond, A2 = cyclo(C3-C9)alkane or cyclo(C5-C9)alkene] or a pro-drug thereof, or a pharmaceutically acceptable salt thereof, which are useful as medicament. Biol. data for compd. II is given.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 1998:631744 CAPLUS
DN 129:310895
TI Benzamide compounds and their use as neovascularization inhibitors
IN Inaba, Takayuki; Tada, Hiroki; Iwamura, Hiroyuki

L6 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
AN 1998:631744 CAPLUS
DN 129:310895
TI Benzamide compounds and their use as neovascularization inhibitors
IN Inaba, Takayuki; Tada, Hiroki; Iwamura, Hiroyuki

PA Japan Tobacco, Inc., Japan
 SO Jpn. Kokai Tokkyo Koho, 106 pp.
 CODEN: JKXXAF

DT Patent
 LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10259176	A2	19980929	JP 1997-84463	19970317
				JP 1997-84463	19970317


OS MARPAT 129:310895

IT 214846-51-2P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (Benzamide compds. and their use as neovascularization inhibitors)

RN 214846-51-2 CAPLUS


CN 2H-1,4-Benzoxazine-7-carboxamide, 3,4-dihydro-3-oxo-4-(phenylmethyl)-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



I



II

AB The inhibitors contain benzamides I [R1 = H, NO₂, halo, cyano, lower alkoxy, NR₁₁R₁₂ (R₁₁, R₁₂ = H, acyl); R₂ = H, NO₂, halo, OR₁₃ (R₁₃ = lower alkyl, aralkyl, cycloalkyl); R₃ = X₃(CH₂)_mR₁₄ [R₁₄ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, (un)substituted lower alkyl, cycloalkyl, acyl, alkenyl, H; X₃ = O, NHCO, OSO₂, NR₁₇ (R₁₇ = H, lower alkyl); m = 0-5], II (R₁₅, R₁₆ = H, lower alkoxy, amino, lower alkyl, CO₂H, OH); R₂ and R₃ may be bonded to form a condensed 1,3-oxazole ring; R₄ = H, OR₁₉ (R₁₉ = lower alkyl, aralkyl, cycloalkyl); R₃ and R₄ may be bonded to form a condensed 1,3-oxazole, 1,4-oxazine, or pyrimidine ring; R₅ = H, NO₂, alkenyl, NHR₂₈ (R₂₈ = H, acyl, lower alkoxy carbonyl); R₆ = H, (un)substituted lower alkyl; R₅ and R₆ may be bonded to form a condensed pyrimidine, diazepine, or pyridine ring; R₇ = H, lower alkoxy; R₈ = X₄(CH₂)_tR₃₀ [X₄ = O, CH₂, CO, or NR₂₈; R₃₀ = H, lower alkyl, aralkyl, direct bond], t = CONH, OSO₂, SO₂NH, NR₃₁ (R₃₁ = H, lower alkyl, aralkyl), (un)substituted 0-5; R₃₀ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, H, OH, halo, lower alkyl, lower alkoxy, cycloalkyl, acyl, cyano, CO₂R₃₂ (R₃₂ = H, lower alkyl); R₉ = H, lower alkoxy carbonyl, halo, OR₃₃ (R₃₃ = H, lower alkyl, aralkyl), CONHR₃₄ (R₃₄ = H, lower alkyl, aralkyl); R₇ and R₈, R₈ and R₉ may be bonded to form a 1,3-oxazole ring; X₁, X₂ = X, N; dotted line represents an optional double bond]. I are useful for treatment of rheumatoid arthritis, diabetic retinopathy, neoplasms, etc. IC₅₀ of 4-benzyloxy-N-(4-benzyloxyphenyl)-3-methoxybenzamide (prepn. given) against bFGF- or VEGF-induced proliferation of HUVEC was 0.85 .mu.M.

AB The inhibitors contain benzamides I [R1 = H, NO₂, halo, cyano, lower alkoxy, NR₁₁R₁₂ (R₁₁, R₁₂ = H, acyl); R₂ = H, NO₂, halo, OR₁₃ (R₁₃ = lower alkyl, aralkyl, cycloalkyl); R₃ = X₃(CH₂)_mR₁₄ [R₁₄ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, (un)substituted lower alkyl, cycloalkyl, acyl, alkenyl, H; X₃ = O, NHCO, OSO₂, NR₁₇ (R₁₇ = H, lower alkyl); m = 0-5], II (R₁₅, R₁₆ = H, lower alkoxy, amino, lower alkyl, CO₂H, OH); R₂ and R₃ may be bonded to form a condensed 1,3-oxazole ring; R₄ = H, OR₁₉ (R₁₉ = lower alkyl, aralkyl, cycloalkyl); R₃ and R₄ may be bonded to form a condensed 1,3-oxazole, 1,4-oxazine, or pyrimidine ring; R₅ = H, NO₂, alkenyl, NHR₂₈ (R₂₈ = H, acyl, lower alkoxy carbonyl); R₆ = H, (un)substituted lower alkyl; R₅ and R₆ may be bonded to form a condensed pyrimidine, diazepine, or pyridine ring; R₇ = H, lower alkoxy; R₈ = X₄(CH₂)_tR₃₀ [X₄ = O, CH₂, CO, CONH, OSO₂, SO₂NH, NR₃₁ (R₃₁ = H, lower alkyl, aralkyl), direct bond], t = 0-5; R₃₀ = (un)substituted Ph, (un)substituted heteroaryl, (un)substituted amino, H, OH, halo, lower alkyl, lower alkoxy, cycloalkyl, acyl, cyano, CO₂R₃₂ (R₃₂ = H, lower alkyl); R₉ = H, lower alkoxy carbonyl, halo, OR₃₃ (R₃₃ = H, lower alkyl, aralkyl), CONHR₃₄ (R₃₄ = H, lower alkyl, aralkyl); R₇ and R₈, R₈ and R₉ may be bonded to form a 1,3-oxazole ring; X₁, X₂ = X, N; dotted line represents an optional double bond]. I are useful for treatment of rheumatoid arthritis, diabetic retinopathy, neoplasms, etc. IC₅₀ of 4-benzyloxy-N-(4-benzyloxyphenyl)-3-methoxybenzamide (prepn. given) against bFGF- or VEGF-induced proliferation of HUVEC was 0.85 .mu.M.

L6 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1992:255611 CAPLUS
 DN 116:255611
 TI Preparation of oxazolyl derivatives
 IN Janssens, Frans Eduard; Sommen, Francois Maria; Dierckx, Ann Christina Joannes; Cooymans, Ludwig Paul
 PA Janssen Pharmaceutica N. V., Belg.
 SO PCT Int. Appl., 63 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9201687	A1	19920206	WO 1991-EP1291	19910709
	W: AU, BB, BG, BR, CA, FI, HU, JP, KP, KR, LK, MC, MG, NO, PL, RO, SU RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FR, GA, GB, GN, GR, IT, LU, ML, MR, NL, SE, SN, TD, TG			US 1990-554326 A	19900719
	US 5217980	A	19930608	US 1991-723862	19910701
				US 1990-554326 B2	19900719
	AU 9182141	A1	19920218	AU 1991-82141	19910709
	AU 644202	B2	19931202		
				US 1990-554326 A	19900719
	EP 539421	A1	19930505	WO 1991-EP1291 A	19910709
	EP 539421	B1	19980923	EP 1991-912700	19910709
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE			US 1990-554326 A	19900719
				WO 1991-EP1291 W	19910709
	JP 05508839	T2	19931209	JP 1991-511644	19910709
	JP 3070951	B2	20000731		

HU 64340	A2	19931228	US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709 HU 1993-97 19910709
RU 2059636	C1	19960510	US 1990-554326 A 19900719 RU 1992-16607 19910709 US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709
PL 169361	B1	19960731	PL 1991-297611 19910709 US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709
RO 111768	B1	19970130	RO 1946-93000 19910709 US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709
RO 111768	B1	19970130	RO 1993-46 19910709 US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709
AT 171449	E	19981015	AT 1991-912700 19910709 US 1990-554326 A 19900719
ES 2121784	T3	19981216	ES 1991-912700 19910709 US 1990-554326 A 19900719
IL 98864	A1	19951208	IL 1991-98864 19910717 US 1990-554326 A 19900719
ZA 9105653	A	19930331	ZA 1991-5653 19910718 US 1990-554326 A 19900719
CZ 279344	B6	19950412	CZ 1991-2240 19910718 US 1990-554326 A 19900719
SK 278133	B6	19960207	SK 1991-2240 19910718 US 1990-554326 A 19900719
CN 1058215	A	19920129	CN 1991-104902 19910719
CN 1043640	B	19990616	US 1990-554326 A 19900719
NO 9300156	A	19930118	NO 1993-156 19930118 US 1990-554326 A 19900719 WO 1991-EP1291 W 19910709
US 5278165	A	19940111	US 1993-35854 19930323 US 1990-554326 B219900719 US 1991-723862 A319910701


OS MARPAT 116:255611

IT 141567-66-0P 141567-90-0P 141567-91-1P
 141568-16-3P 141568-38-9P 141568-88-9P
 141568-95-8P 141569-03-1P 141569-11-1P
 141569-15-5P 141569-44-0P 141569-52-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (prepn. of, as antiallergic)


RN 141567-66-0 CAPLUS

CN 2-Oxazolemethanol, 5-[[2-[[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]amino]-1H-benzimidazol-1-yl]methyl] - (9CI) (CA INDEX NAME)




RN 141567-90-0 CAPLUS

CN Phenol, 4-[2-[4-[[1-[(2-methoxyphenyl)ethyl]-1H-benzimidazol-2-yl]amino]ethyl]- (9CI) (CA INDEX NAME)



RN 141567-91-1 CAPLUS


CN Phenol, 4-[2-[4-[[1-[(2,4-dimethoxyphenyl)ethyl]-1H-benzimidazol-2-yl]amino]ethyl]-, trihydrobromide (9CI) (CA INDEX NAME)



● 3 HBr


RN 141568-16-3 CAPLUS

CN 1H-Benzimidazol-2-amine, N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-1-[(2-methoxyphenyl)ethyl]- (9CI) (CA INDEX NAME)




RN 141568-38-9 CAPLUS

CN 1H-Benzimidazol-2-amine, 1-[(2-methyl-5-oxazolyl)methyl]-N-[1-[(3,4,5-trimethoxyphenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 141568-88-9 CAPLUS

CN 3H-Imidazo[4,5-b]pyridin-2-amine, N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-3-[(2-methyl-5-oxazolyl)methyl]- (9CI) (CA INDEX NAME)




RN 141568-95-8 CAPLUS

CN 1H-Imidazo[4,5-c]pyridin-2-amine, N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-1-[(2-methyl-5-oxazolyl)methyl]- (9CI) (CA INDEX NAME)




RN 141569-03-1 CAPLUS

CN 9H-Purin-8-amine, N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-9-[(2-methyl-5-oxazolyl)methyl]- (9CI) (CA INDEX NAME)



RN 141569-11-1 CAPLUS

CN Phenol, 4-[2-[4-[[3-[(2-methyl-5-oxazolyl)methyl]-3H-imidazo[4,5-b]pyridin-2-yl]amino]-1-piperidinyl]ethyl]- (9CI) (CA INDEX NAME)




RN 141569-15-5 CAPLUS

CN 3H-Imidazo[4,5-c]pyridin-2-amine, N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-3-[(2-methyl-5-oxazolyl)methyl]-, (2E)-2-butenedioate (2:3) (9CI) (CA INDEX NAME)

CM 1

CRN 141569-14-4



CMF C25 H30 N6 O2



CM 2


CRN 110-17-8
CMF C4 H4 O4

Double bond geometry as shown.

RN 141569-44-0 CAPLUS
CN 1H-Benzimidazol-2-amine, 1-[(2,4-dimethyl-5-oxazolyl)methyl]-N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)RN 141569-52-0 CAPLUS
CN 1H-Benzimidazol-2-amine, 1-[(2,5-dimethyl-4-oxazolyl)methyl]-N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]-, (2E)-2-butenedioate (1:2) (9CI) (CA INDEX NAME)

CM 1


CRN 141569-51-9
CMF C27 H33 N5 O2




CM 2

CRN 110-17-8
CMF C4 H4 O4

Double bond geometry as shown.



GI




AB Title compds. I (A1-A4 = CH:CHCH:CH, N:CHCH:CH, CH:NCH:CH, CH:CHN:CH, CH:CHCH:N, N:CHN:CH, CH:NCH:N wherein 1 or 2 H may be replaced by halo, C1-6 alkyl, C1-6 alkoxy, HO, F3C; R = H, C1-4 alkyl; R1 = H, C1-6 alkyl, HO-C1-6-alkyl; D = C1-4 alkanediyl; B = R2N wherein R2 = H, C1-4 alkyl, H2C, O, S, SO, SO2; L = H, C1-12 alkyl, C3-6 cycloalkyl, (aryl) C3-6 alkenyl, C1-6 alkylcarbonyl, C1-6 alkoxy carbonyl, arylcarbonyl, etc.; m = 1,2; n = 0-2), stereoisomer or salt thereof, useful as antiallergic (no data), are prep'd. 5-(Bromomethyl)-2-methyloxazole, Et, 4-[(1H-benzimidazol-2-yl)amino]-1-piperidinecarboxylate, Na2CO3 and DMF were stirred for 18 h at 80.degree. to give after work-up I [A1-A4 = CH:CHCH:CH, R = H, (R1)m = 2-Me, D = H2C, B = NH, L = EtO2C, n = 1].


=> d 19 fbib hitstr abs total

L9 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:515125 CAPLUS
 DN 137:210415
 TI Discovery and structure-activity relationship of N-(ureidoalkyl)-benzyl-piperidines as potent small molecule CC chemokine receptor-3 (CCR3) antagonists
 AU De Lucca, George V.; Kim, Ui T.; Johnson, Curt; Vargo, Brian J.; Welch, Patricia K.; Covington, Maryanne; Davies, Paul; Solomon, Kimberly A.; Newton, Robert C.; Trainor, George L.; Decicco, Carl P.; Ko, Soo S.
 CS Experimental Station, Bristol-Myers Squibb Company, Wilmington, DE, 19880-0336, USA
 SO Journal of Medicinal Chemistry (2002), 45(17), 3794-3804
 CODEN: JMCMAR; ISSN: 0022-2623
 PB American Chemical Society
 DT Journal
 LA English
 IT 275810-47-4P 275810-58-7P
 RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (discovery and structure-activity relationship of N-(ureidoalkyl)benzylpiperidines as CCR3 receptors antagonists)
 RN 275810-47-4 CAPLUS
 CN Urea, N-[1-(phenylmethyl)-4-piperidinyl]-N'-(2-[(4-(phenylmethyl)-1-piperidinyl)methyl]phenyl)- (9CI) (CA INDEX NAME)



RN 275810-58-7 CAPLUS
 CN Urea, N-[1-(phenylmethyl)-4-piperidinyl]-N'-(2-[(4-(phenylmethyl)-1-piperidinyl)methyl]phenyl)- (9CI) (CA INDEX NAME)




AB Structure-activity relationship (SAR) studies of initial screening hits from our corporate library of compds. and a structurally related series of CCR1 receptor antagonists were used to det. that an N-(alkyl)benzylpiperidine is an essential pharmacophore for selective CCR3 antagonists. Further SAR studies that introduced N-(ureidoalkyl) substituents improved the binding potency of these compds. from the micromolar to the low nanomolar range. This new series of compds. also displays highly potent, in vitro functional CCR3-mediated antagonism of eotaxin-induced Ca²⁺ mobilization and chemotaxis of human eosinophils.

RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d l11 fbib hitstr abs total

L11 ANSWER 1 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:335076 CAPLUS
 DN 138:353831
 TI Preparation of 2-carboxypyrrroles as tyrosine kinase inhibitors
 IN Trotter, B. Wesley; Bell, Ian M.; Zartman, C. Blair; Lindsley, Craig;
 Zhao, Zhijian
 PA Merck & Co., Inc., USA
 SO PCT Int. Appl., 208 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003035615	A2	20030501	WO 2002-US33920	20021021
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
				US 2001-343119PP	20011025
OS	MARPAT	138:353831			
IT	518067-63-5P, 2-tert-Butoxycarbonyl-4-methoxycarbonyl-5-[(1-benzylpiperidin-4-yl)amino]methyl]-3-ethyl-1H-pyrrole 518067-64-6P , 2-tert-Butoxycarbonyl-4-methoxycarbonyl-5-[(1-benzylpiperidin-4-yl)amino]methyl]-3-ethyl-1H-pyrrole trifluoroacetate				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(tyrosine kinase inhibitor; prepn. of carboxypyrrroles as tyrosine kinase inhibitors for treatment cancer, diabetes, autoimmune disorders, hyperproliferation disorders, aging, acromegaly, and Crohn's disease)				
RN	518067-63-5 CAPLUS				
CN	1H-Pyrrole-2,4-dicarboxylic acid, 3-ethyl-5-[(1-(phenylmethyl)-4-piperidinyl)amino]methyl-, 2-(1,1-dimethylethyl) 4-methyl ester (9CI) (CA INDEX NAME)				




RN 518067-64-6 CAPLUS

CN 1H-Pyrrole-2,4-dicarboxylic acid, 3-ethyl-5-[[[1-(phenylmethyl)-4-piperidinyl]amino]methyl]-, 2-(1,1-dimethylethyl) 4-methyl ester, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

CRN 518067-63-5


CMF C26 H37 N3 O4





CM 2

CRN 76-05-1

CMF C2 H F3 O2



GI



AB Title compds. I [wherein V = (cyclo)alkyl, aryl, heterocyclyl, or CO; Ra and Rb = independently H, OR7, or (un)substituted alkyl, aryl, or heterocyclyl; R1 = independently H, halo, OR7, COR7, CO2R7, CON(R6)2, N(R7)2, SO2N(R5)2, or (un)substituted (cyclo)alkyl, aryl, or heterocyclyl; R2 = CO2R7, (CRb2)nN(R7)2, CON(R7)2, CONR7OR7, CONH(CRb2)qR7, CONR7NHCOR7, CONR7SO2OR7, (CRb2)nOR7, CONH(CRb2)qCON(R7)2, or (un)substituted alkyl or aryl; R3 and R7 = independently H or (un)substituted alkyl, aralkyl, aryl, or heterocyclyl(alkyl); R4 = (un)substituted alkyl, aryl, aralkyl, or heterocyclyl; R5 = independently H or (un)substituted alkyl, aryl, or heterocyclyl; R6 = independently H, OR7, or (un)substituted alkyl, aralkyl, aryl, or heterocyclyl(alkyl); m = 0-6; n = 0-6; p = 0-6; q = 0-5; and pharmaceutically acceptable salts or stereoisomers thereof] were prep'd. for inhibiting, modulating, and/or regulating signal transduction of both receptor type and non-receptor type tyrosine kinases. For example, addn. of PhCH₂COCl to Meldrum's acid and subsequent treatment with t-BuOH gave tert-Bu 3-oxo-4-phenylbutanoate (no data). Cyclization with NaNO₂ and Et 3-oxobutanoate in the presence of Zn and NH₄OAc, followed by oxidn. and reductive addn. of 4-chloroaniline provided II. Compds. of the invention inhibited insulin-like growth factor I receptor (IGF-1R) or insulin receptor (IR) kinase activity with IC₅₀ values of <100 μM. Thus, I are useful for the treatment of protein kinase related disorders, such as cancer, diabetes, autoimmune disorders, hyperproliferation disorders, aging, acromegaly, and Crohn's disease (no data).

L11 ANSWER 2 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:331563 CAPLUS


DN 139:32012

TI Predicting the Genotoxicity of Secondary and Aromatic Amines Using Data Subsetting To Generate a Model Ensemble

AU Mattioni, Brian E.; Kauffman, Gregory W.; Jurs, Peter C.; Custer, Laura L.; Durham, Stephen K.; Pearl, Greg M.

CS Department of Chemistry, The Pennsylvania State University, University

Park, PA, 16802, USA
 SO Journal of Chemical Information and Computer Sciences (2003), 43(3),
 949-963
 CODEN: JCISD8; ISSN: 0095-2338
 PB American Chemical Society
 DT Journal
 LA English
 IT 68844-77-9, Astemizole
 RL: ADV (Adverse effect, including toxicity); PRP (Properties); BIOL
 (Biological study)
 (predicting genotoxicity of secondary and arom. amines using genetic
 algorithm search engine for data subsetting to generate model ensembles
 based on various mol. descriptors)
 RN 68844-77-9 CAPLUS
 CN 1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-
 methoxyphenyl)ethyl]-4-piperidinyl] - (9CI) (CA INDEX NAME)



AB Binary quant. structure-activity relationship (QSAR) models are developed to classify a data set of 334 arom. and secondary amine compds. as genotoxic or nongenotoxic based on information calcd. solely from chem. structure. Genotoxic endpoints for each compd. were detd. using the SOS Chromotest in both the presence and absence of an S9 rat liver homogenate. Compds. were considered genotoxic if assay results indicated a pos. genotoxicity hit for either the S9 inactivated or S9 activated assay. Each compd. in the data set was encoded through the calcn. of numerical descriptors that describe various aspects of chem. structure (e.g. topol., geometric, electronic, polar surface area). Furthermore, five addnl. descriptors that focused on the secondary and arom. nitrogen atoms in each mol. were calcd. specifically for this study. Descriptor subsets were examd. using a genetic algorithm search engine interfaced with a k-Nearest Neighbor fitness evaluator to find the most information-rich subsets, which ultimately served as the final predictive models. Models were chosen for their ability to minimize the total no. of misclassifications, with special attention given to those models that possessed fewer occurrences of pos. toxicity hits being misclassified as nontoxic (false negatives). In addn., a subsetting procedure was used to form an ensemble of models using different combinations of compds. in the training and prediction sets. This was done to ensure that consistent results could be obtained regardless of training set compn. The procedure also allowed for each compd. to be externally validated three times by different training set data with the resultant predictions being used in a "majority rules" voting scheme to produce a consensus prediction for each member of the data set. The individual models produced an av. training set classification rate of 71.6% and an av. prediction set classification rate of 67.7%. However, the model ensemble was able to correctly classify the genotoxicity of 72.2% of all prediction set compds.

RE.CNT 83 THERE ARE 83 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:98039 CAPLUS
 DN 138:153534
 TI Preparation of benzimidazolyl-substituted quinolinone derivatives and analogs, with inhibitory action against vascular endothelial growth factor receptor tyrosine kinase, and useful as anticancer agents
 IN Renhowe, Paul A.; Pecchi, Sabina; Machajewski, Timothy D.; Shafer, Cynthia M.; Taylor, Clarke; McCrea, William R.; McBride, Christopher; Jazan, Elisa
 PA Chiron Corporation, USA
 SO U.S. Pat. Appl. Publ., 69 pp., Cont.-in-part of U.S. Pat. Appl. 2002 107,392.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003028018	A1	20030206	US 2002-116117	20020405
				US 2000-232159PP	20000911
				US 2001-951265	A220010911
	US 2002107392	A1	20020808	US 2001-951265	20010911
	US 6605617	B2	20030812		
	US 2003158224	A1	20030821	US 2000-232159PP	20000911
				US 2002-284017	20021030
				US 2000-232159PP	20000911
				US 2001-951265	A120010911

PATENT FAMILY INFORMATION:

FAN 2002:220574

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022598	A1	20020321	WO 2001-US42131	20010911
	WO 2002022598	C1	20021121		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2000-232159PP	20000911
	AU 2001093275	A5	20020326	AU 2001-93275	20010911
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
	EP 1317442	A1	20030611	EP 2001-973722	20010911
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
	NO 2003001097	A	20030325	NO 2003-1097	20030310
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911

OS MARPAT 138:153534
 IT 405170-47-0P, 6-Chloro-3-(5-(morpholin-4-yl)-1H-benzimidazol-2-yl)-


4-[[1-(phenylmethyl)piperidin-4-yl]amino]quinolin-2(1H)-one
405170-62-9P, 6-Chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[[1-(phenylmethyl)piperidin-4-yl]amino]quinolin-2(1H)-one

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; prepn. of benzimidazolyl-substituted quinolinone derivs. and analogs as VEGFR tyrosine kinase-inhibiting anticancer agents)


RN 405170-47-0 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[[1-(phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



RN 405170-62-9 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-methyl-1-piperazinyl)-1H-benzimidazol-2-yl]-4-[[1-(phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. of formulas I and II are provided [for I: Z = O, S, (un)substituted NH; Y = certain OH derivs., CHO, esters and amides of CO₂H, certain NH₂ derivs.; R₁-R₄ = H, halo, cyano, NO₂, OH or derivs., NH₂ or derivs., (un)substituted amidinyl, guanidinyl, alk(en/yn)yl, aryl, heterocyclyl, CHO, CO₂H and esters and amides; R₅-R₈ = H, halo, NO₂, OH or derivs., NH₂ or derivs., SH or derivs., cyano, etc.; R₉ = H, OH, (un)substituted alkoxy or aryloxy, NH₂ or derivs., (un)substituted alkyl or aryl, CHO, alkanoyl, aroyl; for II: A, B, D, E = C or N, with at least one being N; Y = H, OH or derivs., SH or derivs., NH₂ or derivs., cyano, various acyl groups, (un)substituted alk(en/yn)yl, aralkyl, heterocycloalkyl, aryl, etc.; R₁-R₈ = H, halo, NO₂, cyano, OH or derivs., NH₂ or derivs., acyl, SH or derivs., etc.; R₉ = H, OH, (un)substituted alkoxy, aryloxy, NH₂ or derivs., aryl, CHO, alkanoyl, aroyl]. Also provided are pharmaceutical formulations including the compds. or their pharmaceutically acceptable salts and a pharmaceutically acceptable carrier, which may be prep'd. by mixing the compds. or salts with a carrier and water. A disclosed method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient. Claims include tautomers of the compds., pharmaceutically acceptable salts, and pharmaceutically acceptable salts of the tautomers. I and II are inhibitors of receptor tyrosine kinases, and particularly of vascular endothelial growth factor receptor (VEGFR) tyrosine kinase. As such, they are inhibitors of angiogenesis, and thereby act as anticancer agents. Approx 270 invention compds. are listed, with detailed preps. given for about 50 compds. Several general preparatory methods are discussed in detail. For instance, cyclocondensation of Et 2-(benzimidazol-2-yl)acetate with the corresponding ortho-amino nitrile (preps. given), carried out in refluxing ClCH₂CH₂Cl in the presence of SnCl₄, gave the invention quinolinone III. Many compds. I and II had in vitro IC₅₀ values of less than 10 .mu.M with respect to flt-1 (VEGFR1), KDR (VEGFR2) and bFGF kinases (recombinant, expressed in Sf9 insect cells).

L11 ANSWER 4 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:906175 CAPLUS

DN 138:14074

TI Preparation of benzo[g]quinoxalines for use against infectious diseases

IN Pato, Janos; Keri, Gyoergy; Oerfi, Laszlo; Waczek, Frigyes; Horvath, Zoltan; Banhegyi, Peter; Szabadkai, Istvan; Marosfalvi, Jenoe; Hegymegi-barakonyi, Balint; Szekelyhidi, Zsolt; Greff, Zoltan; Choidas, Axel; Bacher, Gerald; Daub, Henrik; Obert, Sabine; Kurtenbach, Alexander; Habenberger, Peter

PA Axxima Pharmaceuticals Ag, Germany; et al.

SO PCT Int. Appl., 237 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE	
PI	WO 2002094796	A2	20021128	WO 2002-EP5573	20020521	
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,				


TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 EP 2001-112289 A 20010518
 US 2001-292325PP 20010522
 US 2001-298902PP 20010619
 EP 2001-115508 A 20010627


OS MARPAT 138:14074

IT 476637-76-0P, N,N'-Bis(1-benzylpiperidin-4-yl)benzo[g]quinoxaline-
 2,3-diamineRL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)(drug candidate; prepn. of benzo[g]quinoxalines for use against
 infectious diseases)

RN 476637-76-0 CAPLUS

CN Benzo[g]quinoxaline-2,3-diamine, N,N'-bis[1-(phenylmethyl)-4-piperidinyl]-
 (9CI) (CA INDEX NAME)

GI



AB The present invention relates to benzo[g]quinoxaline derivs. (shown as I; e.g. 2,3-bis(2-thienyl)benzo[g]quinoxaline and benzo[g]quinoxalin-2-yl(3-bromophenyl)amine), processes for manufg. said benzo[g]quinoxaline derivs., the use of the benzo[g]quinoxaline derivs. as pharmaceutically active agents, esp. for the prophylaxis and/or treatment of infectious diseases and opportunistic infections, diabetes, cancer, inflammation, as well as compns. contg. at least one benzo[g]quinoxaline deriv. and/or pharmaceutically acceptable salt thereof. Further, the present invention is directed to methods for preventing and/or treating of infectious diseases, diabetes, cancer, and inflammation using the inventive benzo[g]quinoxaline derivs. The inventive benzo[g]quinoxaline derivs.


exert their antiproliferative effect on *M. bovis* BCG and *M. tuberculosis* Erdmann at concns. between <<1 .mu.M and 32 .mu.M. In contrast, growth of *E. coli* XI-1 blue was not affected by benzo[g]quinoxaline derivs. at concns. >10 .mu.M. The benzo[g]quinoxaline compds. are able to inhibit HI virus replication up to 63% after 6 days at a concn. of 1 .mu.M. 5,10-Dibromo-2-(thiophen-3-yl)-3-(thiophen-2-yl)benzo[g]quinoxaline is able to decrease the activity of the herpes viral target UL-97 by 75%. Results for inhibition of HCMV target RICK for 5 I, of influenza replication for 7 I, of hepatitis B virus for 5 I, of TNF.alpha. signaling for 11 I, of human cellular protein kinases (Akt, Abl, PDGFR, Src) for 7 I, of A549 and Jurkat cells for 18 I, of human cellular protein kinase Akt known as a target for diabetes for 4 I, and of human protein kinases SRPK1 and SRPK2 (indicative of hepatitis B virus replication inhibition) for 8 and 1 I, resp., are tabulated. Results for activation of the insulin receptor InsR by 3 I, effect of 2 I on viability of Huh-5-2 replicon cells by the Alamar Blue toxicity assay, effect of 2 I on autonomous replication of hepatitis C virus replicons in the Huh-5-2 cell line by luciferase reporter assay, are tabulated. In I: R1 and R2 = -(CH₂)p-NH-(CH₂)n-R9, -(CH₂)s-S-(CH₂)m-R10, -(CH₂)m-O-(CH₂)p-R11, -(CH₂)r-R3, -CH:CH-R11, -(CH₂)m-CH(OH)(CH₂)p-R11, -(CH₂)q-R11, -R9, R10, -R12, -R13, etc. R3, R4, R5, R6, R7, and R8 = -H, -F, -Cl, -Br, -I, -SO₃H, -SO₃NH₂, -(CH₂)s-COOR16, -(CH₂)p-COOR17, -OR16, -SR16, -NR16R17, -OOCR16, -OOCR17, -NH-CO-R16, -NH-CO-R17, -CO-NH-R16, -CO-NH-R17, -NO₂, -N₃, -CN, -OCN, -NCO, -SCN, -NCS, CO-R16, CO-R17, -COCN, -CONR16R17, -SOR16, -SO₂R16, -SO₂R17, -SO₃R16, -SO₃R17, OCF₃. R9, R10, and R11 = -CN, NR16R17, -NHR16, NHR17, etc. R12, R13, R14, and R15 = R3, R4, R5, R6, R16, R17, CH(CO₂R16)(CO₂R17), CH(CN)(CO₂R16), CH(CN)C(O)NHAr (Ar = R₁₄- and R₁₅-substituted phenyl); R16 and R17 = -H, -CH₃, -C₂H₅, -Pr, -CHMe₂, -Bu, -C₅H₁₁, -C₆H₁₃, -cyclo-C₆H₁₁, -cyclo-C₅H₉, -cyclo-C₄H₇, -cyclo-C₃H₅, -(CH₂)r-CHMe₂, -CHMeEt, -CMe₃, -CH:CH₂, -CH₂-CH:CH₂, Ph, --CH₂Ph, -C₂H₄Ph, -CH(CN)₂, -CF₃, -CCl₃, -CBr₃, -C₂F₅, -(CH₂)r-OH, -CH₂F, -CH₂Cl, -CH₂Br, -CH₂I, -CHF₂, -CHCl₂, -CHBr₂, -(CH₂)r-SH, -C₆H₄-CH₃, -C₆H₃Me₂, pyridyl, 2-pyrimidinyl, etc. M = 0-6, n = 0-6, p = 0-6, q = 0-6, r = 1-6, s = 0-6. Also claimed are the corresponding N-oxides in position 1 and/or 4 of these compds., the corresponding reduced forms of these compds. wherein the double bond in position 1 and/or 3 is hydrogenated, and pharmaceutically acceptable salts of I. About 42 example preps. and 406 compds. with characterization data are included. 1H-benzo[g]quinoxaline-2-one was prep'd. in 90% yield by dissolving 20 mmol 2,3-diaminonaphthalene in a mixt. of 5 mL DMF and 50 mL EtOH and adding 5 mL aq. soln. (50%) of glyoxalic acid and the mixt. was stirred for 2 h at reflux temp. The reaction mixt. was cooled to room temp. and the product was filtered, washed two times with Et₂O and dried.

L11 ANSWER 5 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:888714 CAPLUS
 DN 137:384765
 TI Preparation of novel 4-anilinoquinoline-3-carboxamides as JAK3 kinase inhibitors
 IN Larsson, Joakim; Sjoe, Peter
 PA Astrazeneca AB, Swed.
 SO PCT Int. Appl., 97 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----

PI WO 2002092571 A1 20021121 WO 2002-SE875 20020506
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 SE 2001-1675 A 20010511

OS MARPAT 137:384765
 IT **476190-02-0P**, 7-[(1-Benzyl-4-piperidinyl)amino]-6-methoxy-4-(2-methylphenylamino)-3-quinolinecarboxamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of novel 4-anilinoquinoline-3-carboxamides as JAK3 kinase inhibitors)
 RN 476190-02-0 CAPLUS
 CN 3-Quinolinecarboxamide, 6-methoxy-4-[(2-methylphenyl)amino]-7-[(1-phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



GI




AB The title compds. [I; n = 0-1; X = NR3, O; Ar = (un)substituted Ph, indolyl, pyrazolyl, etc.; R = H, alkyl; R1, R2 = H, halo, NO2, etc.; or R1 and R2 are linked together as OCH2O or OCH2CH2O] which are JAK3 kinase inhibitors, useful in treating asthma, host vs. graft rejection/transplantation or rheumatoid arthritis, were prep'd. E.g., a 7-step synthesis of I [X = NH; n = 0; Ar = 3-(hydroxymethyl)-2-

methylphenyl; R1 = OCH₂Ph; R2 = OMe}, starting from 4-nitroguaiacol potassium salt, was given. The exemplified compds. I showed IC₅₀ of < 25 μM in JAK3 HTRF assay.


RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 6 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:594844 CAPLUS
 DN 137:140518
 TI Preparation of thiazolyl-, oxazolyl-, pyrrolyl-, and imidazolyl- acid amide derivatives as inhibitors of phosphodiesterase IV isozymes
 IN Marfat, Anthony; McKechney, Michael William
 PA Pfizer Products Inc., USA
 SO PCT Int. Appl., 249 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002060898	A1	20020808	WO 2001-IB2728	20011224
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2001-265486PP	20010131
US	2002123520	A1	20020905	US 2002-62145	20020131
US	6559168	B2	20030506		
US	2003130254	A1	20030710	US 2001-265486PP	20010131
				US 2002-300959	20021120
				US 2001-265486PP	20010131
				US 2002-62145	A320020131
US	2003186974	A1	20031002	US 2002-300950	20021120
				US 2001-265486PP	20010131
				US 2002-62145	A320020131
OS	MARPAT 137:140518				
IT	68844-77-9 , Atemizole				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(combination therapy with PDE4 inhibitors; prepn. of thiazolyl-, oxazolyl-, pyrrolyl-, and imidazolyl- acid amide derivs. as inhibitors of PDE4 isoenzymes)				
RN	68844-77-9 CAPLUS				
CN	1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)				



GI



I



II

AB Title compds. I [wherein p = 0-1; q = 0-1; provided that when q = 0, n = 2; m = 0-3; n = 1-2; W1 and W2 = independently O, SOO-2, or NR3; or W2 = (un)substituted methylene; Y = SOO-2, O, NOO-1, NR3, or (un)substituted methylene; ; RA and RB = independently H, F, CF3, alkyl, or (un)substituted cycloalkyl, Ph, or benzyl; or when m = 1, CRARB = (un)substituted spiro; RC and RD have the same meaning as RA and RB except that one of them must be H; R1 and R2 = H, F, Cl, CN, NO2, (fluoro)alkyl, alkynyl, alkoxy, phenoxy, carbamoyl, etc.; R3 = H, alkyl, Ph, benzyl, alkoxy, phenoxy, etc.; R4, R5, and R6 = H, F, Cl, and (un)substituted (cyclo)alkyl, alkenyl, alkynyl, Ph, benzyl, pyridyl, alkoxy, phenoxy, acyl, carboxy, CN, NO2, carbamoyl, ureido, (hetero)aryl, etc.; G1 and G2 = independently (un)satd. carbocyclyl or heterocyclyl; E = (un)substituted carboxy, carbamoyl, acyl, hydroxyalkyl, cyanoalkyl, acylamino, ureido, amino, heterocyclyl, etc.] were prep'd. as inhibitors of PDE4 (no data). For example, 4-(3-cyanophenoxy)thiazole-5-carboxylic acid was treated with

2-(4-aminomethylphenyl)propan-2-ol in the presence of EDC1 and HOBT in DMF to give the thiazolamide II. I are useful in the treatment of diseases regulated by the activation and degranulation of eosinophils, esp. asthma, chronic bronchitis, and chronic obstructive pulmonary disease (no data). In addn., I may be used in combination therapy with a wide variety of other therapeutic agents.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 7 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:293652 CAPLUS
 DN 136:325531
 TI Preparation of (poly)azanaphthalenyl carboxamides as HIV integrase inhibitors
 IN Anthony, Neville J.; Gomez, Robert P.; Young, Steven D.; Egbertson, Melissa; Wai, John S.; Zhuang, Linghang; Embrey, Mark; Tran, Lekhanh; Melamed, Jeffrey Y.; Langford, H. Marie; Guare, James P.; Fisher, Thorsten E.; Jolly, Samson M.; Kuo, Michelle S.; Perlow, Debra S.; Bennett, Jennifer J.; Funk, Timothy W.
 PA Merck & Co., Inc., USA
 SO PCT Int. Appl., 434 pp.
 CODEN: PIXXD2

DT Patent
 LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002030930	A2	20020418	WO 2001-US31456	20011009
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2000-239707PP	20001012
				US 2001-281656PP	20010405
AU	2002011527	A5	20020422	AU 2002-11527	20011009
				US 2000-239707PP	20001012
				US 2001-281656PP	20010405
				WO 2001-US31456W	20011009
EP	1326865	A2	20030716	EP 2001-979582	20011009
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-239707PP	20001012
				US 2001-281656PP	20010405
				WO 2001-US31456W	20011009
US	2003055071	A1	20030320	US 2001-973853	20011010
				US 2000-239707PP	20001012
				US 2001-281656PP	20010405

PATENT FAMILY INFORMATION:

FAN 2002:293653

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002030931	A2	20020418	WO 2001-US42564	20011009
	WO 2002030931	A3	20021024		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS,
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT,
 RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
 UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2002011874	A5	20020422	US 2000-239707PP 20001012
			US 2001-281656PP 20010405
EE 200300145	A	20030616	AU 2002-11874 20011009
			US 2000-239707PP 20001012
US 2003055071	A1	20030320	US 2001-281656PP 20010405
			WO 2001-US42564W 20011009
NO 2003001672	A	20030605	EE 2003-145 20011009
			US 2000-239707PP 20001012
			US 2001-281656PP 20010405
			WO 2001-US42564W 20011009


OS MARPAT 136:325531

IT 410543-58-7P, 5-[(1-Benzylpiperidin-4-yl)amino]-N-(3,5-dichlorobenzyl)-8-hydroxy-1,6-naphthyridine-7-carboxamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)



(HIV integrase inhibitor; prepn. of (poly)azanaphthalenyl carboxamides as HIV integrase inhibitors for treatment of AIDS)

RN 410543-58-7 CAPLUS

CN 1,6-Naphthyridine-7-carboxamide, N-[(3,5-dichlorophenyl)methyl]-8-hydroxy-5-[[1-(phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



GI



AB Title compds., including certain quinoline carboxamide and naphthyridine carboxamide derivs., I [wherein A = (un)substituted Ph or Ph fused to a carbocycle; L = a single bond, or (un)substituted alkyl, alkenyl, alkylcycloalkylalkyl, or alkyl-M-alkyl; M = NR_a, OCO, or CO₂; X = N or CQ₁; Y = N or CQ₂, provided that X and Y are not both N; Z₁ = N or CQ₃; Z₂ = N or CQ₄; Z₃ = N or CH; Q₁-Q₄ = independently H, halo, CN, NR₁CR₁O, or (un)substituted alkyl, alkoxy, alkenyl, alkynyl, carbamoyl, carboximidamido, amino, etc.; or C₂Q₂Q₃ = (un)substituted 5- or 6-membered carbocycle or heterocycle; R₁ and R₂ = independently H, OH, halo, NO₂, CN, or (un)substituted alkyl, alkenyl, alkoxy, amino, sulfonylamino, etc.; R₃ and R₄ = independently H, halo, CN, NO₂, OH, alkenyl, or (un)substituted alkyl, amino, sulfonylamino, etc.; R₅ = H, CN, CN, or (un)substituted alkyl or aryl; Ra = independently H or (halo)alkyl; or pharmaceutically acceptable salts thereof] were prep'd. I are inhibitors of HIV integrase and inhibitors of HIV replication, and are useful in the prevention or treatment of infection by HIV and the treatment of AIDS, as compds. or pharmaceutically acceptable salts, or as ingredients in pharmaceutical compns., optionally in combination with other antivirals, immunomodulators, antibiotics, or vaccines. For example, Mitsunobu reaction of iso-Pr 3-(hydroxymethyl)pyridine-2-carboxylate with Me N-[(4-methylphenyl)sulfonyl]glycinate, followed by cyclization in the presence on NaOMe, afforded Me 8-hydroxy-1,6-naphthyridine-7-carboxylate. Coupling with 3,5-dichlorobenzylamine in toluene gave II. Representative compds. were assayed for the inhibition of acute HIV infection of T-lymphoid cells and demonstrated IC₉₅ values of < 20 .mu.M.

L11 ANSWER 8 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:240760 CAPLUS

DN 136:279470

TI Preparation of 6-[(substituted phenyl)methyl]quinoline and quinazoline derivatives as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases

IN Angibaud, Patrick Rene; Venet, Marc Gaston; Saha, Ashis Kumar; Mevellec, Laurence Anne

PA Janssen Pharmaceutica N.V., Belg.

SO PCT Int. Appl., 97 pp.


CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002024683	A1	20020328	WO 2001-EP10895	20010918
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	
				EP 2000-203366 A	20000925
	AU 2001093829	A5	20020402	AU 2001-93829	20010918
				EP 2000-203366 A	20000925
				WO 2001-EP10895W	20010918
	EP 1322636	A1	20030702	EP 2001-974276	20010918
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			EP 2000-203366 A	20000925
				WO 2001-EP10895W	20010918
OS	MARPAT 136:279470				
IT	406164-50-9P				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(farnesyl transferase inhibitor; prepn. of quinoline and quinazoline derivs. as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases)				
RN	406164-50-9 CAPLUS				
CN	Tetrazolo[1,5-a]quinazoline-7-methanol, 5-(3-chlorophenyl)-.alpha.-[(1-methyl-1H-imidazol-5-yl)-.alpha.-[4-[[[1-(phenylmethyl)-4-piperidinyl]amino]methyl]phenyl]- (9CI) (CA INDEX NAME)				



GI



AB Title compds. I [wherein m and n = independently 0-5; q = 0-3; Y₁Y₂ = C:N, C:CR₉, CHNR₉, or CHCHR₉; C₉ = H, halo, CN, (cyclo)alkyl, hydroxyalkyl, alkoxy(alkyl), aminoalkyl, (amino)alkenyl, (amino)alkynyl, halocarbonyl, hydroxycarbonyl, alkoxycarbonyl, aryl, (un)substituted amino or carbamoyl, etc.; R₁ and R₂ = independently azido, OH, halo, CN, NO₂, trihalomethyl, alkoxy, aryloxy, heterocyclyloxy, alkylthio, or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, carbamoyl, amino, sulfamoyl, etc.; or 2 adjacent R₁ = OCH₂O, OCH₂CH₂O, OCH:CH, OCH₂CH₂, OCH₂CH₂CH₂, CH:CHCH:CH; R₃ = H, halo, CN, alkenyl, alkynyl, hydroxycarbonyl, alkoxycarbonyl, aryl, heterocyclyl, alkoxy, alkylthio, (un)substituted (cyclo)alkyl or amino, etc.; R₄ = (un)substituted imidazolyl, triazolyl, or pyridyl; R₅ = CN, OH, halo, alkenyl, alkynyl, hydroxycarbonyl, alkoxycarbonyl, or (un)substituted (cyclo)alkyl, alkoxy, amino, or carbamoyl, etc.; R₆ = halo or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, alkylthio, carboxy, carbamoyl, acyl(amino), etc.; R₇ = O or S; or R₆R₇ = (un)substituted CH:CHN:, CH:NN:, N:NN:, N:CHN:, CH:CHCH:, CH:NCH:, CONHCH:, N:NCH:, or CH₂(CH₂)₀₋₁CH₂N: ; or pharmaceutically acceptable salts, N-oxides, or stereochem. isomeric forms thereof] were prepd. For example, 6-bromo-2-chloro-4-(3-chlorophenyl)quinoline (6-step prepn. given) was coupled with 4-(diethoxymethyl)benzaldehyde in the presence of BuLi in THF to give the 6-quinolinemethanol (64%), which was treated with MnO₂ in 1,4-dioxane to afford the methanone. Methoxylation using MeONa in MeOH (74%), followed by addn. of 1-methyl-1H-imidazole in the presence of BULi and ClSiEt₃ in THF, gave 4-(3-chlorophenyl)-.alpha.-[4-(diethoxymethyl)phenyl]-2-methoxy-.alpha.-(1-methyl-1H-imidazol-5-yl)-6-quinolinemethanol (56%). The latter was refluxed in HCl for 24 h, cooled, poured out into H₂O, and stirred at room temp. for 1 h to afford the quinolinone II.bul.HCl (98%). I have potent farnesyl transferase inhibitory effect and are useful for inhibiting proliferative diseases and growth of tumors expressing an activated ras oncogene (no data).

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 9 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2002:220574 CAPLUS
DN 136:263158
TI Benzimidazolyl-substituted quinolinone derivatives and analogs, with inhibitory action against vascular endothelial growth factor receptor tyrosine kinase, and useful as anticancer agents
IN Renhowe, Paul; Pecchi, Sabina; Machajewski, Tim; Shafer, Cynthia; Taylor, Clarke; McCrea, Bill; McBride, Chris; Jazan, Elisa; Wernette-Hammond, Mary-Ellen; Harris, Alex
PA Chiron Corporation, USA

SO PCT Int. Appl., 207 pp.
CODEN: PIXXD2

DT Patent
LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022598	A1	20020321	WO 2001-US42131	20010911
	WO 2002022598	C1	20021121		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2000-232159PP	20000911
	AU 2001093275	A5	20020326	AU 2001-93275	20010911
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
EP	1317442	A1	20030611	EP 2001-973722	20010911
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
	NO 2003001097	A	20030325	NO 2003-1097	20030310
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911

PATENT FAMILY INFORMATION:

FAN 2003:98039

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003028018	A1	20030206	US 2002-116117	20020405
				US 2000-232159PP	20000911
				US 2001-951265 A2	20010911
	US 2002107392	A1	20020808	US 2001-951265	20010911
	US 6605617	B2	20030812		
				US 2000-232159PP	20000911
	US 2003158224	A1	20030821	US 2002-284017	20021030
				US 2000-232159PP	20000911
				US 2001-951265 A1	20010911

OS MARPAT 136:263158


IT 405170-47-0P, 6-Chloro-3-(5-(morpholin-4-yl)-1H-benzimidazol-2-yl)-4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)-one
405170-62-9P, 6-Chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)-one

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; prepn. of benzimidazolyl-substituted quinolinone derivs. and analogs as VEGFR tyrosine kinase-inhibiting anticancer agents)


RN 405170-47-0 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)-4-piperidinyl)amino]- (9CI) (CA INDEX NAME)



RN 405170-62-9 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-methyl-1-piperazinyl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)-4-piperidinyl)amino]- (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. of formulas I and II are provided [for I: Z = O, S, (un)substituted NH; Y = certain OH derivs., CHO, esters and amides of CO₂H, certain NH₂ derivs.; R1-R4 = H, halo, cyano, NO₂, OH or derivs., NH₂ or derivs., (un)substituted amidinyl, guanidinyl, alk(en/yn)yl, aryl, heterocyclyl, CHO, CO₂H and esters and amides; R5-R8 = H, halo, NO₂, OH or derivs., NH₂ or derivs., SH or derivs., cyano, etc.; R9 = H, OH, (un)substituted alkoxy or aryloxy, NH₂ or derivs., (un)substituted alkyl or aryl, CHO, alkanoyl, aroyl; for II: A, B, D, E = C or N, with at least one being N; Y = H, OH or derivs., SH or derivs., NH₂ or derivs., cyano, various acyl groups, (un)substituted alk(en/yn)yl, aralkyl, heterocycloalkyl, aryl, etc.; R1-R8 = H, halo, NO₂, cyano, OH or derivs., NH₂ or derivs., acyl, SH or derivs., etc.; R9 = H, OH, (un)substituted


alkoxy, aryloxy, NH₂ or derivs., aryl, CHO, alkanoyl, aroyl]. Also provided are pharmaceutical formulations including the compds. or their pharmaceutically acceptable salts and a pharmaceutically acceptable carrier, which may be prep'd. by mixing the compds. or salts with a carrier and water. A disclosed method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient. Claims include tautomers of the compds., pharmaceutically acceptable salts, and pharmaceutically acceptable salts of the tautomers. I and II are inhibitors of receptor tyrosine kinases, and particularly of vascular endothelial growth factor receptor (VEGFR) tyrosine kinase. As such, they are inhibitors of angiogenesis, and thereby act as anticancer agents. Approx 270 invention compds. are listed, with detailed preps. given for about 50 compds. Several general preparatory methods are discussed in detail. For instance, cyclocondensation of Et 2-(benzimidazol-2-yl)acetate with the corresponding ortho-amino nitrile (preps. given), carried out in refluxing ClCH₂CH₂Cl in the presence of SnCl₄, gave the invention quinolinone III. Many compds. I and II had in vitro IC₅₀ values of less than 10 .mu.M with respect to flt-1 (VEGFR1), KDR (VEGFR2) and bFGF kinases (recombinant, expressed in Sf9 insect cells).

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 10 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:123617 CAPLUS
 DN 136:183819
 TI Preparation of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors
 IN Wang, Wei-Bo; Curtin, Michael L.; Fakhoury, Stephen A.; Gwaltney, Stephen L.; Hasvold, Lisa A.; Hutchins, Charles W.; Li, Qun; Lin, Nan-Horng; Nelson, Lissa Taka Jennings; O'Connor, Steve; Sham, Hing L.; Sullivan, Gerard M.; Wang, Gary T.; Wang, Xilu
 PA USA
 SO U.S. Pat. Appl. Publ., 189 pp.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002019527	A1	20020214	US 2001-842391	20010425
				US 2000-200165PP	20000427

OS MARPAT 136:183819
 IT 371761-79-4P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors)
 RN 371761-79-4 CAPLUS
 CN Benzonitrile, 4-[(1-methyl-1H-imidazol-5-yl)[[1-(phenylmethyl)-4-piperidinyl]amino)methyl]-2-(1-naphthalenyl)-, trihydrochloride (9CI) (CA INDEX NAME)



● 3 HCl

GI



AB Title compds. (I) were prep'd. Thus, 2-MeC₆H₄C₆H₃(CN)(CHO)-2,5 was condensed with 1-methyl-2-triethylsilyl-1H-imidazole (prepn. each given) and the product O-arylated to give title compd. II. Data for biol. activity of I were given.

L11 ANSWER 11 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:107318 CAPLUS

DN 136:151163

TI Preparation of indazole derivatives as JNK enzyme inhibitors

IN Bhagwat, Shripad S.; Satoh, Yoshitaka; Sakata, Steven T.

PA Signal Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 412 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002010137	A2	20020207	WO 2001-US23890	20010730
	WO 2002010137	C2	20030206		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO,					

RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2000-221799PP 20000731

US 2002103229 A1 20020801 US 2001-910950 20010723

US 2000-221799PP 20000731

EP 1313711 A2 20030528 EP 2001-957332 20010730

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

US 2000-221799PP 20000731

WO 2001-US23890W 20010730

OS MARPAT 136:151163


IT 395107-63-8P, N-[1-Benzyl-4-piperidyl]-3-[5-(1H-1,2,4-triazol-3-yl)-1H-indazol-3-yl]benzamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of indazole derivs. as JNK enzyme inhibitors)

RN 395107-63-8 CAPLUS

CN Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]-3-[5-(1H-1,2,4-triazol-3-yl)-1H-indazol-3-yl] - (9CI) (CA INDEX NAME)




AB Indazole derivs., 3-R1A-5-R2-1H-indazoles (1), having activity as selective inhibitors of JNK are disclosed. In 1: A is a direct bond, -(CH₂)_a, -(CH₂)_bCH:CH(CH₂)_c, or -(CH₂)_bC.tplbond.C(CH₂)_c; R1 is aryl, heteroaryl or heterocycle fused to Ph, each being optionally substituted with 1-4 R3; R2 is -R₃, -R₄, -(CH₂)_bC(O)R₅, -(CH₂)_bC(:O)OR₅, -(CH₂)_bC(O)NR₅R₆, -(CH₂)_bC(O)NR₅(CH₂)_cC(O)R₆, -(CH₂)_bNR₅C(O)R₆, -(CH₂)_bNR₅C(O)NR₆R₇, -(CH₂)_bNR₅R₆, -(CH₂)_bOR₅, -(CH₂)_bSOdR₅ or -(CH₂)_bSO₂NR₅R₆. A is 1-6; b and c are the same or different and are 0-4; d is 0-2. R3 is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, heterocycle, substituted heterocycle, heterocyclealkyl, substituted heterocyclealkyl, -C(O)OR₈, -C(O)R₈, -C(O)NR₈R₉, -C(O)NR₈OR₉, -SO₂NR₈R₉, -NR₈SO₂R₉, -CN, -NO₂, -NR₈R₉, -NR₈C(O)R₉, -NR₈C(O)(CH₂)_bOR₉, -NR₈C(O)(CH₂)_bR₉, -O(CH₂)_bNR₅R₉, or heterocycle fused to Ph. R4 is alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, each being optionally substituted with 1-4 R3, or R4 is halogen or hydroxy. R5, R6 and R7 are the same or different and are H, alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, wherein each of R5, R6 and R7 are optionally substituted with 1-4 R3. R8 and R9 are the same or different and at each occurrence independently H, alkyl, aryl, arylalkyl, heterocycle, or heterocyclealkyl, or R8 and R9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R8, R9, and R8 and R9 taken together to form a heterocycle

are optionally substituted with 1-4 R3 with the proviso that: when A is a direct bond and R1 is Ph, R2 is not Me, methoxy, C(O)CH₃ or C(O)H; when A is a direct bond and R1 is 4-Me-Ph, R2 is not Me; when A is a direct bond and R1 is 4-F-Ph, R2 is not trifluoromethyl; when A is a direct bond or -C.tpbond.C- and R1 is Ph, R2 is not -COOEt; and when A is a direct bond and R1 is 6,7-dimethoxyisoquinolin-1-yl, R2 is not hydroxy. Such compds. have utility in the treatment of a wide range of conditions that are responsive to JNK inhibition. Thus, methods of treating such conditions are also disclosed, as are pharmaceutical compns. contg. one or more compds. of the above compds. Many of the claimed compds. have IC₅₀ values $\text{IC}_{50} \text{req. } 0.5 \text{ } \mu\text{M}$ in the JNK2 assay, e.g. 5-[3-(4-fluorophenyl)-1H-indazol-5-yl]-2H-1,2,3,4-tetrazole. Although the methods of prepn. are not claimed, >400 example prepn.s. are included.

L11 ANSWER 12 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2001:798200 CAPLUS
 DN 135:344482
 TI Preparation of substituted 4-(heteroaryl methyl)benzonitriles as farnesyltransferase inhibitors
 IN Wang, Wei-Bo; Curtin, Michael L.; Fakhoury, Stephen A.; Gwaltney, Stephen L., II; Hasvold, Lisa A.; Hutchins, Charles W.; Li, Qui; Lin, Nan-Horng; Jennings Nelson, Lissa Taka; O'Connor, Stephen J.; Sham, Hing L.; Sullivan, Gerald M.; Wang, Gary T.; Wang, Xilu
 PA Abbott Laboratories, USA
 SO PCT Int. Appl., 305 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001081316	A2	20011101	WO 2001-US13678	20010425
	WO 2001081316	A3	20020523		
		W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		
		RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG		US 2000-563256 A 20000427
					US 2001-822205 A 20010402
EP	1276726	A2	20030122	EP 2001-932712	20010425
		R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		US 2000-563256 A 20000427
					US 2001-822205 A 20010402
					WO 2001-US13678W 20010425
OS	MARPAT 135:344482				
IT	371761-79-4P				
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(prepn. of substituted 4-(heteroaryl methyl)benzonitriles as farnesyltransferase inhibitors)				
RN	371761-79-4 CAPLUS				

CN Benzonitrile, 4-[(1-methyl-1H-imidazol-5-yl)[[1-(phenylmethyl)-4-piperidinyl]amino]methyl]-2-(1-naphthalenyl)-, trihydrochloride (9CI) (CA INDEX NAME)



● 3 HCl

GI



AB The title compds. [I; A1 = (un)substituted alkylene, etc.; R1 = halo, cycloalkyl, aryl, heteroaryl; R2 = heteroaryl selected from imidazolyl, pyrazolyl, pyrrolyl, etc.] and their pharmaceutically acceptable salts which farnesyltransferase, were prep'd. E.g., 3-step synthesis of the benzonitrile II.HCl which 88% inhibition of farnesyltransferase at 10⁻⁶ M, was given.

L11 ANSWER 13 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2001:31473 CAPLUS

DN 134:100864

TI Indazole compounds and pharmaceutical compositions for inhibiting protein kinases, and methods for their use

IN Kania, Robert Steven; Bender, Steven Lee; Borchardt, Allen J.; Braganza,

John F.; Cripps, Stephan James; Hua, Ye; Johnson, Michael David; Johnson, Theodore Otto, Jr.; Luu, Hiep The; Palmer, Cynthia Louise; Reich, Siegfried Heinz; Tempczyk-russell, Anna Maria; Teng, Min; Thomas, Christine; Varney, Michael David; Wallace, Michael Brennan

PA Agouron Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 439 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001002369	A2	20010111	WO 2000-US18263	20000630
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				US 1999-142130PP 19990702
BR	2000012352	A	20020514	BR 2000-12352	20000630
				US 1999-142130PP 19990702	
				WO 2000-US18263W 20000630	
EP	1218348	A2	20020703	EP 2000-943375	20000630
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
				US 1999-142130PP 19990702	
				WO 2000-US18263W 20000630	
JP	2003503481	T2	20030128	JP 2001-507809	20000630
				US 1999-142130PP 19990702	
				WO 2000-US18263W 20000630	
US	6531491	B1	20030311	US 2001-983786	20011025
				US 1999-142130PP 19990702	
				US 2000-609335 B320000630	
US	6534524	B1	20030318	US 2001-983783	20011025
				US 1999-142130PP 19990702	
				US 2000-609335 B320000630	
NO	2001005797	A	20020301	NO 2001-5797	20011128
				US 1999-142130PP 19990702	
				WO 2000-US18263W 20000630	
ZA	2001010061	A	20030206	ZA 2001-10061	20011206
				US 1999-142130PP 19990702	
BG	106380	A	20020930	BG 2002-106380	20020201
				US 1999-142130PP 19990702	
				WO 2000-US18263W 20000630	

OS MARPAT 134:100864

IT 319466-31-4P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of combinatorial libraries of aryl-substituted indazole derivs.
 as modulators and inhibitors of protein kinases in the treatment of
 tumor growth, cellular proliferation, and angiogenesis)

RN 319466-31-4 CAPLUS


CN Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]-2-[[3-[(1E)-2-(2-

pyridinyl)ethenyl]-1H-indazol-6-yl]thio]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.



GI



AB Indazole compds. I [R1 = substituted or unsubstituted aryl or heteroaryl, R₃CH:CH, R₃N:CH; R2 = substituted or unsubstituted aryl, heteroaryl, Y-X; R3 = substituted or unsubstituted alkyl alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl; Y = O, S, C(:CH₂), CO, SO, SO₂, alkylidene, NH, N(C₁-C₈ alkyl); X = substituted or unsubstituted aryl, heteroaryl, NH(alkyl), NH(cycloalkyl), NH(heterocycloalkyl), NH(aryl), NH(heteroaryl), NH(alkoxy), NH(dialkylamide)] and their pharmaceutically acceptable prodrugs, active metabolites, and salts are disclosed. The compds. modulate and/or inhibit the activity of certain protein kinases. In particular, I and pharmaceutical compns. contg. them are capable of mediating tyrosine kinase signal transduction, and thereby modulate and/or inhibit unwanted cell proliferation. The invention is also directed to the therapeutic or prophylactic use of pharmaceutical compns. contg. such compds., and to methods of treating cancer and other disease states assoc'd. with unwanted angiogenesis and/or cellular proliferation, such as diabetic retinopathy, neovascular glaucoma, rheumatoid arthritis, and psoriasis, by administering effective amts. of such compds. E.g., I [R1 = (E)-3,4-(MeO)2C₆H₃CH:CH; R2 = 4-HO-3-MeOC₆H₃] (II) was prep'd. from 6-aminoindazole by diazotization and substitution with iodide, protection of the indazole nitrogen with 2,4,6-Me₃C₆H₂SO₂Cl, coupling of the regioisomeric mixt. with 4-(methoxymethoxy)-3-methoxybenzeneboronic acid in the presence of dichlorobis(triphenylphosphine)palladium, and deprotection of the indazole moiety and iodination at the 3-position of the indazole. Treatment of the 3-indazolyl iodide with sec-butyllithium, phenyllithium, and DMF, regioselective protection of the indazole with 2,4,6-Me₃C₆H₂SO₂Cl, olefination with 3,4-dimethoxybenzyltriphenylphosphonium bromide, deprotection of the indazole, deprotection of the methoxymethyl group, and equilibration of the double bond with iodine gave II. Biol. data on protein kinase inhibition, cell proliferation


inhibition, neovascularization inhibition, and i.p. and oral bioavailability, are given.

L11 ANSWER 14 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1995:772553 CAPLUS
 DN 123:199300
 TI Preparation of diaminopurinylibofuranuronamide derivatives as antiinflammatories.
 IN Gregson, Michael; Ayres, Barry Edward; Ewan, George Blanch; Ellis, Frank; Knight, John
 PA Glaxo Group Ltd., UK
 SO PCT Int. Appl., 112 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9417090	A1	19940804	WO 1994-EP145	19940118
	W: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			GB 1993-1000	A 19930120
CA	2153688	AA	19940804	CA 1994-2153688	19940118
				GB 1993-1000	A 19930120
AU	9458851	A1	19940815	AU 1994-58851	19940118
AU	679714	B2	19970710	GB 1993-1000	A 19930120
				WO 1994-EP145	W 19940118
ZA	9400335	A	19941024	ZA 1994-335	19940118
				GB 1993-1000	A 19930120
EP	680488	A1	19951108	EP 1994-905100	19940118
EP	680488	B1	19980408	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE	
				GB 1993-1000	A 19930120
				WO 1994-EP145	W 19940118
CN	1119440	A	19960327	CN 1994-191527	19940118
CN	1043997	B	19990707	GB 1993-1000	A 19930120
				JP 1994-516652	19940118
JP	08505864	T2	19960625	GB 1993-1000	A 19930120
				WO 1994-EP145	W 19940118
AT	164849	E	19980415	AT 1994-905100	19940118
				GB 1993-1000	A 19930120
ES	2117249	T3	19980801	ES 1994-905100	19940118
				GB 1993-1000	A 19930120
RU	2129561	C1	19990427	RU 1995-122754	19940118
				GB 1993-1000	A 19930120
				WO 1994-EP145	W 19940118
SK	281229	B6	20010118	SK 1995-918	19940118
				GB 1993-1000	A 19930120
				WO 1994-EP145	W 19940118
IL	108372	A1	19980615	IL 1994-108372	19940119
				GB 1993-1000	A 19930120
FI	9503489	A	19950913	FI 1995-3489	19950719
				GB 1993-1000	A 19930120


NO 9502872	A	19950913	WO 1994-EP145	W 19940118
			NO 1995-2872	19950719
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
US 5925624	A	19990720	US 1995-446727	19950918
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
US 5889178	A	19990330	US 1997-934540	19970922
			GB 1993-1000	A 19930120
			US 1995-446727	A319950918
OS	MARPAT 123:199300			
IT	167297-77-0P			
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)			
	(prepn. of diaminopurinylribofuranuronamide derivs. as antiinflammatories)			
RN	167297-77-0 CAPLUS			
CN	.beta.-D-Ribofuranuronamide, 1-deoxy-1-[6-[(2,2-diphenylethyl)amino]-2-[(1-(phenylmethyl)-4-piperidinyl]amino]-9H-purin-9-yl]-N-ethyl- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.




IT	167297-68-9P
	RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
	(prepn. of diaminopurinylribofuranuronamide derivs. as antiinflammatories)
RN	167297-68-9 CAPLUS
CN	.beta.-D-Ribofuranuronamide, 1-deoxy-1-[6-[(2,2-diphenylethyl)amino]-2-[(1-(phenylmethyl)-4-piperidinyl]amino]-9H-purin-9-yl]-N-ethyl-2,3-O-(1-methylethylidene)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



GI



AB Title compds. [I; R₁ = H, C₃-8 cycloalkyl, C₁-6 alkyl; R₂ = (substituted) C₃-8 cycloalkyl, C₃-8 cycloalkyl-C₁-6 alkyl, pyrrolidin-3-yl, 2-oxopyrrolidin-4-yl, 2-oxopyrrolidin-5-yl, piperidin-3-yl, piperidin-4-yl, etc.; Q = O, S], were prep'd. Title compds. are useful as antiinflammatory agents, particularly in the treatment of patients with inflammatory conditions who are susceptible to leukocyte-induced tissue damage. Thus, (trans)-1-[2-[(4-aminocyclohexyl)amino]-6-[(2,2-diphenylethyl)amino]-9H-purin-9-yl]-1-deoxy-N-ethyl-2,3-O-(1-methylethylidene)-.beta.-D-ribofuranuronamide was stirred with aq. CF₃CO₂H to give (trans)-1-[2-[(4-aminocyclohexyl)amino]-6-[(2,2-diphenylethyl)amino]-9H-purin-9-yl]-1-deoxy-N-ethyl-.beta.-D-ribofuranonamide. The latter was 25 times more potent than NECA for inhibiting O₂- generation from neutrophils stimulated with fMLP, and inhibited ovalbumin-induced eosinophil accumulation in sensitized guinea pigs with ED₅₀ = 10 .mu.g/kg i.p.

L11 ANSWER 15 OF 15 CAPLUS COPYRIGHT 2003 ACS on STN

AN 1989:574103 CAPLUS

DN 111:174103

TI Preparation of piperidine-containing heterocycles as analgesics and anesthetics

IN Lin, Bor Sheng; Scheblein, Joseph W.

PA BOC Inc., USA
 SO U.S., 28 pp.
 CODEN: USXXAM
 DT Patent
 LA English
 FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4831192	A	19890516	US 1987-139896	19871231
	EP 328830	A1	19890823	EP 1988-312149	19881221
	EP 328830	B1	19940601		
	R: BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE			US 1987-139896	19871231
				US 1987-139899	19871231
	ES 2054836	T3	19940816	ES 1988-312149	19881221
				US 1987-139896	19871231
				US 1987-139899	19871231
	JP 01301676	A2	19891205	JP 1988-332751	19881228
				US 1987-139896	19871231

PATENT FAMILY INFORMATION:

FAN 1989:423391


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4791120	A	19881213	US 1987-139899	19871231
	US 4871749	A	19891003	US 1988-256792	19881011
				US 1987-139899	19871231
	AU 8826604	A1	19890713	AU 1988-26604	19881206
	AU 616708	B2	19911107		
	NO 8805463	A	19890703	US 1987-139899	19871231
	NO 174553	B	19940214	NO 1988-5463	19881208
	NO 174553	C	19940525		
	IL 88645	A1	19930708	US 1987-139899	19871231
				IL 1988-88645	19881209
	EP 328830	A1	19890823	US 1987-139899	19871231
	EP 328830	B1	19940601	EP 1988-312149	19881221
	R: BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE			US 1987-139896	19871231
				US 1987-139899	19871231
	ES 2054836	T3	19940816	ES 1988-312149	19881221
				US 1987-139896	19871231
				US 1987-139899	19871231
	JP 01213278	A2	19890828	JP 1988-332752	19881228
				US 1987-139899	19871231
	DK 8807328	A	19890701	DK 1988-7328	19881230
				US 1987-139899	19871231
	FI 8806057	A	19890701	FI 1988-6057	19881230
	FI 92065	B	19940615		
	FI 92065	C	19940926		
	CN 1035285	A	19890906	US 1987-139899	19871231
	CN 1023010	B	19931208	CN 1989-100056	19881230
				US 1987-139899	19871231
OS	CASREACT 111:174103; MARPAT 111:174103				
IT	968-86-5P 120070-52-2P 120070-54-4P				
	120070-55-5P 120115-93-7P				
	RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT				

(Reactant or reagent)

(prepn. and reaction of, in prepn. of analgesics and anesthetics)


RN 968-86-5 CAPLUS

CN 4-Piperidinecarbonitrile, 4-(phenylamino)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 120070-52-2 CAPLUS

CN 4-Piperidinamine, 4-(1-methyl-1H-1,2,4-triazol-5-yl)-N-phenyl-1-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 120070-54-4 CAPLUS

CN 4-Piperidinamine, N-phenyl-1-(phenylmethyl)-4-(1H-tetrazol-5-yl)- (9CI) (CA INDEX NAME)




RN 120070-55-5 CAPLUS

CN 4-Piperidinamine, N-(2-fluorophenyl)-1-(phenylmethyl)-4-(1H-tetrazol-5-yl)- (9CI) (CA INDEX NAME)



RN 120115-93-7 CAPLUS

CN 4-Piperidinamine, 4- (4-methyl-2-thiazolyl) -N-phenyl-1- (phenylmethyl) -
(9CI) (CA INDEX NAME)




IT 120070-56-6

RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, in prepn. of analgesics and anesthetics)

RN 120070-56-6 CAPLUS

CN 4-Piperidinecarbonitrile, 4-[(2-fluorophenyl)amino]-1-(phenylmethyl)-
(9CI) (CA INDEX NAME)



GI For diagram(s), see printed CA Issue.

AB Title compds. I [R1 = (substituted) oxadiazolyl, imidazolyl, triazolyl, tetrazolyl, thiazolyl; R2 = (substituted) Ph; R3 = acyl, alkoxy carbonyl; L = alkyl, alkoxy, thiarylalkyl, (substituted) thiazolylalkyl, etc.] are prep'd. from I (R1 = cyano; R3 = H). Treatment of I (R1 = cyano; R2 = Ph; R3 = H; L = PhCH2) (prep'd. from KCN, PhNH2, and N-benzyl-4-piperidone, CAUTION: HCN evolution) with NaN3 in THF in the presence of AlCl3 gave I (R1 = 1H-tetrazol-5-yl), which was refluxed with Ac2O to afford I (R1 = 5-methyl-1,3,4-oxadiazol-2-yl; R2 = Ph; R3 = Ac; L = PhCH2). The oxalate of the latter showed ED50 >5.0 mg/kg in mice in a hot-plate analgesia

test.

=> d his

(FILE 'HOME' ENTERED AT 10:51:12 ON 13 OCT 2003)

FILE 'REGISTRY' ENTERED AT 10:52:17 ON 13 OCT 2003

L1 STRUCTURE UPLOADED
L2 4255 S L1 SSS FULL

FILE 'CAPLUS' ENTERED AT 10:53:24 ON 13 OCT 2003

L3 1572 S L2
L4 13 S L3 AND OXADIAZOLE
L5 98 S L3 AND IMIDAZOLE
L6 6 S L3 AND OXAZOLE
L7 295 S L3 AND ONE
L8 0 S L3 AND CHEMIKINE
L9 1 S L3 AND CHEMOKINE AND CCR1 AND CCR3
L10 20 S L3 AND TRIAZOLE
L11 15 S L3 AND 1,2,4-TRIAZOLE

=> d l10 fbib hitstr abs total

L10 ANSWER 1 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2003:356269 CAPLUS

DN 138:348761

TI Type 4 phosphodiesterase inhibitors and therapeutic uses thereof

IN Eggenweiler, Hans-Michael; Wolf, Michael

PA Merck Patent G.m.b.H., Germany

SO PCT Int. Appl., 122 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE	
PI	WO 2003037349	A1	20030508	WO 2002-EP9596	20020828	
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	

EP 2001-125394 A 20011031

OS MARPAT 138:348761

IT 68844-77-9, Astemizole


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(phosphodiesterase IV inhibitors, therapeutic uses, and use with other agents)

RN 68844-77-9 · CAPLUS

CN 1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-

methoxyphenyl)ethyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)



AB The invention discloses the use of type 4 phosphodiesterase inhibitors (PDE IV inhibitors) to treat diseases, as well as combinations of PDE IV inhibitors with other drugs.

RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 2 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:335076 CAPLUS

DN 138:353831

TI Preparation of 2-carboxypyrrroles as tyrosine kinase inhibitors

IN Trotter, B. Wesley; Bell, Ian M.; Zartman, C. Blair; Lindsley, Craig; Zhao, Zhijian

PA Merck & Co., Inc., USA

SO PCT Int. Appl., 208 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003035615	A2	20030501	WO 2002-US33920	20021021
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2001-343119PP	20011025


OS MARPAT 138:353831

IT 518067-63-5P, 2-tert-Butoxycarbonyl-4-methoxycarbonyl-5-[(1-benzylpiperidin-4-yl)amino]methyl]-3-ethyl-1H-pyrrole 518067-64-6P, 2-tert-Butoxycarbonyl-4-methoxycarbonyl-5-[(1-benzylpiperidin-4-yl)amino]methyl]-3-ethyl-1H-pyrrole trifluoroacetate

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(tyrosine kinase inhibitor; prepn. of carboxypyrrroles as tyrosine kinase inhibitors for treatment cancer, diabetes, autoimmune disorders, hyperproliferation disorders, aging, acromegaly, and Crohn's disease)

RN 518067-63-5 CAPLUS

CN 1H-Pyrrole-2,4-dicarboxylic acid, 3-ethyl-5-[[[1-(phenylmethyl)-4-piperidinyl]amino]methyl]-, 2-(1,1-dimethylethyl) 4-methyl ester (9CI)
(CA INDEX NAME)


RN 518067-64-6 CAPLUS

CN 1H-Pyrrole-2,4-dicarboxylic acid, 3-ethyl-5-[[[1-(phenylmethyl)-4-piperidinyl]amino]methyl]-, 2-(1,1-dimethylethyl) 4-methyl ester, mono(trifluoroacetate) (9CI) (CA INDEX NAME)

CM 1

CRN 518067-63-5


CMF C26 H37 N3 O4





CM 2

CRN 76-05-1

CMF C2 H F3 O2



GI



AB Title compds. I [wherein V = (cyclo)alkyl, aryl, heterocyclyl, or CO; Ra and Rb = independently H, OR7, or (un)substituted alkyl, aryl, or heterocyclyl; R1 = independently H, halo, OR7, COR7, CO2R7, CON(R6)2, N(R7)2, SO2N(R5)2, or (un)substituted (cyclo)alkyl, aryl, or heterocyclyl; R2 = CO2R7, (CRb2)n(R7)2, CON(R7)2, CONR7OR7, CONH(CRb2)qR7, CONR7NHCOR7, CONR7SO2OR7, (CRb2)nOR7, CONH(CRb2)qCON(R7)2, or (un)substituted alkyl or aryl; R3 and R7 = independently H or (un)substituted alkyl, aralkyl, aryl, or heterocyclyl(alkyl); R4 = (un)substituted alkyl, aryl, aralkyl, or heterocyclyl; R5 = independently H or (un)substituted alkyl, aryl, or heterocyclyl; R6 = independently H, OR7, or (un)substituted alkyl, aralkyl, aryl, or heterocyclyl(alkyl); m = 0-6; n = 0-6; p = 0-6; q = 0-5; and pharmaceutically acceptable salts or stereoisomers thereof] were prep'd. for inhibiting, modulating, and/or regulating signal transduction of both receptor type and non-receptor type tyrosine kinases. For example, addn. of PhCH₂COCl to Meldrum's acid and subsequent treatment with t-BuOH gave tert-Bu 3-oxo-4-phenylbutanoate (no data). Cyclization with NaNO₂ and Et 3-oxobutanoate in the presence of Zn and NH₄OAc, followed by oxidn. and reductive addn. of 4-chloroaniline provided II. Compds. of the invention inhibited insulin-like growth factor I receptor (IGF-1R) or insulin receptor (IR) kinase activity with IC₅₀ values of $\leq 100 \mu M$. Thus, I are useful for the treatment of protein kinase related disorders, such as cancer, diabetes, autoimmune disorders, hyperproliferation disorders, aging, acromegaly, and Crohn's disease (no data).

L10 ANSWER 3 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:331563 CAPLUS


DN 139:32012

TI Predicting the Genotoxicity of Secondary and Aromatic Amines Using Data Subsetting To Generate a Model Ensemble

AU Mattioni, Brian E.; Kauffman, Gregory W.; Jurs, Peter C.; Custer, Laura L.; Durham, Stephen K.; Pearl, Greg M.

CS Department of Chemistry, The Pennsylvania State University, University


Park, PA, 16802, USA
 SO Journal of Chemical Information and Computer Sciences (2003), 43(3),
 949-963
 CODEN: JCISD8; ISSN: 0095-2338
 PB American Chemical Society
 DT Journal
 LA English
 IT 68844-77-9, Astemizole
 RL: ADV (Adverse effect, including toxicity); PRP (Properties); BIOL
 (Biological study)
 (predicting genotoxicity of secondary and arom. amines using genetic
 algorithm search engine for data subsetting to generate model ensembles
 based on various mol. descriptors)
 RN 68844-77-9 CAPLUS
 CN 1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-
 methoxyphenyl)ethyl]-4-piperidinyl] - (9CI) (CA INDEX NAME)




AB Binary quant. structure-activity relationship (QSAR) models are developed to classify a data set of 334 arom. and secondary amine compds. as genotoxic or nongenotoxic based on information calcd. solely from chem. structure. Genotoxic endpoints for each compd. were detd. using the SOS Chromotest in both the presence and absence of an S9 rat liver homogenate. Compds. were considered genotoxic if assay results indicated a pos. genotoxicity hit for either the S9 inactivated or S9 activated assay. Each compd. in the data set was encoded through the calcn. of numerical descriptors that describe various aspects of chem. structure (e.g. topol., geometric, electronic, polar surface area). Furthermore, five addnl. descriptors that focused on the secondary and arom. nitrogen atoms in each mol. were calcd. specifically for this study. Descriptor subsets were examd. using a genetic algorithm search engine interfaced with a k-Nearest Neighbor fitness evaluator to find the most information-rich subsets, which ultimately served as the final predictive models. Models were chosen for their ability to minimize the total no. of misclassifications, with special attention given to those models that possessed fewer occurrences of pos. toxicity hits being misclassified as nontoxic (false negatives). In addn., a subsetting procedure was used to form an ensemble of models using different combinations of compds. in the training and prediction sets. This was done to ensure that consistent results could be obtained regardless of training set compn. The procedure also allowed for each compd. to be externally validated three times by different training set data with the resultant predictions being used in a "majority rules" voting scheme to produce a consensus prediction for each member of the data set. The individual models produced an av. training set classification rate of 71.6% and an av. prediction set classification rate of 67.7%. However, the model ensemble was able to correctly classify the genotoxicity of 72.2% of all prediction set compds.

RE.CNT 83 THERE ARE 83 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L10 ANSWER 4 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2003:295477 CAPLUS
 DN 139:69208
 TI Click Linker: Efficient and High-Yielding Synthesis of a New Family of SPOS Resins by 1,3-Dipolar Cycloaddition
 AU Loeber, Stefan; Rodriguez-Loaiza, Pilar; Gmeiner, Peter
 CS Department of Medicinal Chemistry, Emil Fischer Center, Friedrich-Alexander University, Erlangen, D-91052, Germany
 SO Organic Letters (2003), 5(10), 1753-1755
 CODEN: ORLEF7; ISSN: 1523-7060
 PB American Chemical Society
 DT Journal
 LA English
 OS CASREACT 139:69208
 IT 135385-46-5P 552311-89-4P 552311-97-4P
 552312-06-8P 552312-12-6P 552312-16-0P
 552312-21-7P 552312-25-1P
 RL: CPN (Combinatorial preparation); PAC (Pharmacological activity); BIOL (Biological study); CMBI (Combinatorial study); PREP (Preparation) (prep. and dopamine D4 receptor selectivity and binding of a combinatorial library of amides prepd. on solid-phase using a novel methylindolylmethyltriazolyl linker)
 RN 135385-46-5 CAPLUS
 CN 3-Pyridinecarboxamide, N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 552311-89-4 CAPLUS
 CN Benzo[b]thiophene-2-carboxamide, N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 552311-97-4 CAPLUS
 CN 3-Pyridinecarboxamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 552312-06-8 CAPLUS
 CN 2-Quinolinemcarboxamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]- (9CI)
 (CA INDEX NAME)




RN 552312-12-6 CAPLUS
 CN 2-Quinolinemcarboxamide, N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



RN 552312-16-0 CAPLUS
 CN Benzamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]-2,5-dimethoxy- (9CI) (CA INDEX NAME)




RN 552312-21-7 CAPLUS
 CN Benzamide, 2,5-dimethoxy-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)




RN 552312-25-1 CAPLUS

CN Benzo[b]thiophene-2-carboxamide, N-[1-[(4-cyanophenyl)methyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB Functionalized resins for the prepn. of combinatorial libraries are prep'd. using the 1,3-dipolar cycloaddn. of a resin-bound azide with propargyl-substituted aryl aldehydes to yield a variety of resin-bound triazole-contg. aryl aldehydes as the key step. Reductive amination of the resin-bound aldehydes with amines followed by carbodiimide-mediated coupling of the resin-bound amines with carboxylic acids and cleavage of the amides from the resin with trifluoroacetic acid yields amides. This method allows the prepn. of a variety of linkers for solid-phase synthesis of combinatorial libraries and thus allows the linker to be readily optimized for the prepn. of the desired combinatorial library. In the case of an amide library, a 3-methylindolylmethyltriazolyl linker is found to provide the amide products in high purity and yield. Using the 3-methylindolylmethyltriazolyl linker, a library of twenty amides are prep'd. and tested for binding to dopamine D₄ receptors; I (R = 2-quinolyl, 2-benzothienyl) are found to bind selectively to D₄ dopamine receptors in preference to D₂(long), D₂(short), and D₃ dopamine receptors.

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 5 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:98039 CAPLUS
 DN 138:153534
 TI Preparation of benzimidazolyl-substituted quinolinone derivatives and
 analogs, with inhibitory action against vascular endothelial growth factor
 receptor tyrosine kinase, and useful as anticancer agents
 IN Renhowe, Paul A.; Pecchi, Sabina; Machajewski, Timothy D.; Shafer, Cynthia
 M.; Taylor, Clarke; McCrea, William R.; McBride, Christopher; Jazan, Elisa
 PA Chiron Corporation, USA
 SO U.S. Pat. Appl. Publ., 69 pp., Cont.-in-part of U.S. Pat. Appl. 2002
 107,392.
 CODEN: USXXCO

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003028018	A1	20030206	US 2002-116117	20020405
				US 2000-232159PP	20000911
				US 2001-951265	A220010911
	US 2002107392	A1	20020808	US 2001-951265	20010911
	US 6605617	B2	20030812		
	US 2003158224	A1	20030821	US 2000-232159PP	20000911
				US 2002-284017	20021030
				US 2000-232159PP	20000911
				US 2001-951265	A120010911

PATENT FAMILY INFORMATION:

FAN 2002:220574


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022598	A1	20020321	WO 2001-US42131	20010911
	WO 2002022598	C1	20021121		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2000-232159PP	20000911
	AU 2001093275	A5	20020326	AU 2001-93275	20010911
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
EP	1317442	A1	20030611	EP 2001-973722	20010911
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
NO	2003001097	A	20030325	NO 2003-1097	20030310
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
OS	MARPAT 138:153534				
IT	405170-47-0P, 6-Chloro-3-(5-(morpholin-4-yl)-1H-benzimidazol-2-yl)- 4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)-one 405170-62-9P, 6-Chloro-3-[5-(4-methylpiperazin-1-yl)-1H- benzimidazol-2-yl]-4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)- one				

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; prepn. of benzimidazolyl-substituted quinolinone derivs. and analogs as VEGFR tyrosine kinase-inhibiting anticancer agents)


RN 405170-47-0 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)-4-piperidinyl)amino]- (9CI) (CA INDEX NAME)



RN 405170-62-9 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-methyl-1-piperazinyl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)-4-piperidinyl)amino]- (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. of formulas I and II are provided [for I: Z = O, S, (un)substituted NH; Y = certain OH derivs., CHO, esters and amides of CO2H, certain NH2 derivs.; R1-R4 = H, halo, cyano, NO2, OH or derivs., NH2]


or derivs., (un)substituted amidinyl, guanidinyl, alk(en/yn)yl, aryl, heterocyclyl, CHO, CO₂H and esters and amides; R₅-R₈ = H, halo, NO₂, OH or derivs., NH₂ or derivs., SH or derivs., cyano, etc.; R₉ = H, OH, (un)substituted alkoxy or aryloxy, NH₂ or derivs., (un)substituted alkyl or aryl, CHO, alkanoyl, aroyl; for II: A, B, D, E = C or N, with at least one being N; Y = H, OH or derivs., SH or derivs., NH₂ or derivs., cyano, various acyl groups, (un)substituted alk(en/yn)yl, aralkyl, heterocycloalkyl, aryl, etc.; R₁-R₈ = H, halo, NO₂, cyano, OH or derivs., NH₂ or derivs., acyl, SH or derivs., etc.; R₉ = H, OH, (un)substituted alkoxy, aryloxy, NH₂ or derivs., aryl, CHO, alkanoyl, aroyl]. Also provided are pharmaceutical formulations including the compds. or their pharmaceutically acceptable salts and a pharmaceutically acceptable carrier, which may be prep'd. by mixing the compds. or salts with a carrier and water. A disclosed method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient. Claims include tautomers of the compds., pharmaceutically acceptable salts, and pharmaceutically acceptable salts of the tautomers. I and II are inhibitors of receptor tyrosine kinases, and particularly of vascular endothelial growth factor receptor (VEGFR) tyrosine kinase. As such, they are inhibitors of angiogenesis, and thereby act as anticancer agents. Approx 270 invention compds. are listed, with detailed preps. given for about 50 compds. Several general preparatory methods are discussed in detail. For instance, cyclocondensation of Et 2-(benzimidazol-2-yl)acetate with the corresponding ortho-amino nitrile (preps. given), carried out in refluxing ClCH₂CH₂Cl in the presence of SnCl₄, gave the invention quinolinone III. Many compds. I and II had in vitro IC₅₀ values of less than 10 .mu.M with respect to flt-1 (VEGFR1), KDR (VEGFR2) and bFGF kinases (recombinant, expressed in Sf9 insect cells).

L10 ANSWER 6 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:906175 CAPLUS
 DN 138:14074
 TI Preparation of benzo[*g*]quinoxalines for use against infectious diseases
 IN Pato, Janos; Keri, Gyoergy; Oerfi, Laszlo; Waczek, Frigyes; Horvath,
 Zoltan; Banhegyi, Peter; Szabadkai, Istvan; Marosfalvi, Jenoe;
 Hegymegi-barakonyi, Balint; Szekelyhidi, Zsolt; Greff, Zoltan; Choidas,
 Axel; Bacher, Gerald; Daub, Henrik; Obert, Sabine; Kurtenbach, Alexander;
 Habenberger, Peter
 PA Axxima Pharmaceuticals Ag, Germany; et al.
 SO PCT Int. Appl., 237 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002094796	A2	20021128	WO 2002-EP5573	20020521
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG			

EP 2001-112289 A 20010518
 US 2001-292325PP 20010522
 US 2001-298902PP 20010619
 EP 2001-115508 A 20010627

OS MARPAT 138:14074
 IT 476637-76-0P, N,N'-Bis(1-benzylpiperidin-4-yl)benzo[g]quinoxaline-2,3-diamine
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (drug candidate; prepn. of benzo[g]quinoxalines for use against infectious diseases)
 RN 476637-76-0 CAPLUS
 CN Benzo[g]quinoxaline-2,3-diamine, N,N'-bis[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB The present invention relates to benzo[g]quinoxaline derivs. (shown as I; e.g. 2,3-bis(2-thienyl)benzo[g]quinoxaline and benzo[g]quinoxalin-2-yl(3-bromophenyl)amine), processes for manufg. said benzo[g]quinoxaline derivs., the use of the benzo[g]quinoxaline derivs. as pharmaceutically active agents, esp. for the prophylaxis and/or treatment of infectious diseases and opportunistic infections, diabetes, cancer, inflammation, as well as compns. contg. at least one benzo[g]quinoxaline deriv. and/or pharmaceutically acceptable salt thereof. Further, the present invention is directed to methods for preventing and/or treating of infectious diseases, diabetes, cancer, and inflammation using the inventive benzo[g]quinoxaline derivs. The inventive benzo[g]quinoxaline derivs. exert their antiproliferative effect on *M. bovis* BCG and *M. tuberculosis* Erdmann at concns. between <<1 .mu.M and 32 .mu.M. In contrast, growth of *E. coli* XI-1 blue was not affected by benzo[g]quinoxaline derivs. at concns. >10 .mu.M. The benzo[g]quinoxaline compds. are able to inhibit HI

virus replication up to 63% after 6 days at a concn. of 1 .mu.M. 5,10-Dibromo-2-(thiophen-3-yl)-3-(thiophen-2-yl)benzo[g]quinoxaline is able to decrease the activity of the herpes viral target UL-97 by 75%. Results for inhibition of HCMV target RICK for 5 I, of influenza replication for 7 I, of hepatitis B virus for 5 I, of TNF.alpha. signaling for 11 I, of human cellular protein kinases (Akt, Abl, PDGFR, Src) for 7 I, of A549 and Jurkat cells for 18 I, of human cellular protein kinase Akt known as a target for diabetes for 4 I, and of human protein kinases SRPK1 and SRPK2 (indicative of hepatitis B virus replication inhibition) for 8 and 1 I, resp., are tabulated. Results for activation of the insulin receptor InsR by 3 I, effect of 2 I on viability of Huh-5-2 replicon cells by the Alamar Blue toxicity assay, effect of 2 I on autonomous replication of hepatitis C virus replicons in the Huh-5-2 cell line by luciferase reporter assay, are tabulated. In I: R1 and R2 = -(CH₂)p-NH-(CH₂)n-R9, -(CH₂)s-S-(CH₂)m-R10, -(CH₂)m-O-(CH₂)p-R11, -(CH₂)r-R3, -CH:CH-R11, -(CH₂)m-CH(OH)(CH₂)p-R11, -(CH₂)q-R11, -R9, R10, -R12, -R13, etc. R3, R4, R5, R6, R7, and R8 = -H, -F, -Cl, -Br, -I, -SO₃H, -SO₃NH₂, -(CH₂)s-COOR₁₆, -(CH₂)p-COOR₁₇, -OR₁₆, -SR₁₆, -NR₁₆R₁₇, -OOCR₁₆, -OOCR₁₇, -NH-CO-R₁₆, -NH-CO-R₁₇, -CO-NH-R₁₆, -CO-NH-R₁₇, -NO₂, -N₃, -CN, -OCN, -NCO, -SCN, -NCS, CO-R₁₆, CO-R₁₇, -COCN, -CONR₁₆R₁₇, -SOR₁₆, -SO₂R₁₆, -SO₂R₁₇, -SO₃R₁₆, -SO₃R₁₇, OCF₃. R9, R10, and R11 = -CN, NR₁₆R₁₇, -NHR₁₆, NHR₁₇, etc. R12, R13, R14, and R15 = R3, R4, R5, R6, R16, R17, CH(CO₂R₁₆)(CO₂R₁₇), CH(CN)(CO₂R₁₆), CH(CN)C(O)NHAr (Ar = R₁₄- and R₁₅-substituted phenyl); R16 and R17 = -H, -CH₃, -C₂H₅, -Pr, -CHMe₂, -Bu, -C₅H₁₁, -C₆H₁₃, -cyclo-C₆H₁₁, -cyclo-C₅H₉, -cyclo-C₄H₇, -cyclo-C₃H₅, -(CH₂)r-CHMe₂, -CHMeEt, -CMe₃, -CH:CH₂, -CH₂-CH:CH₂, Ph, --CH₂Ph, -C₂H₄Ph, -CH(CN)₂, -CF₃, -CCl₃, -CBr₃, -C₂F₅, -(CH₂)r-OH, -CH₂F, -CH₂Cl, -CH₂Br, -CH₂I, -CHF₂, -CHCl₂, -CHBr₂, -(CH₂)r-SH, -C₆H₄-CH₃, -C₆H₃Me₂, pyridyl, 2-pyrimidinyl, etc. M = 0-6, n = 0-6, p = 0-6, q = 0-6, r = 1-6, s = 0-6. Also claimed are the corresponding N-oxides in position 1 and/or 4 of these compds., the corresponding reduced forms of these compds. wherein the double bond in position 1 and/or 3 is hydrogenated, and pharmaceutically acceptable salts of I. About 42 example preps. and 406 compds. with characterization data are included. 1H-benzo[g]quinoxaline-2-one was prep'd. in 90% yield by dissolving 20 mmol 2,3-diaminonaphthalene in a mixt. of 5 mL DMF and 50 mL EtOH and adding 5 mL aq. soln. (50%) of glyoxalic acid and the mixt. was stirred for 2 h at reflux temp. The reaction mixt. was cooled to room temp. and the product was filtered, washed two times with Et₂O and dried.

L10 ANSWER 7 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:888714 CAPLUS
 DN 137:384765
 TI Preparation of novel 4-anilinoquinoline-3-carboxamides as JAK3 kinase inhibitors
 IN Larsson, Joakim; Sjoe, Peter
 PA Astrazeneca AB, Swed.
 SO PCT Int. Appl., 97 pp.
 CODEN: PIXXD2

DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
PI WO 2002092571	A1	20021121	WO 2002-SE875	20020506
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				

LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
 UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
 TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
 SE 2001-1675 A 20010511

OS MARPAT 137:384765


IT 476190-02-0P, 7-[(1-Benzyl-4-piperidinyl)amino]-6-methoxy-4-(2-methylphenylamino)-3-quinolinecarboxamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(prepn. of novel 4-anilinoquinoline-3-carboxamides as JAK3 kinase inhibitors)

RN 476190-02-0 CAPLUS

CN 3-Quinolinecarboxamide, 6-methoxy-4-[(2-methylphenyl)amino]-7-[(1-phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



GI




AB The title compds. [I; n = 0-1; X = NR3, O; Ar = (un)substituted Ph, indolyl, pyrazolyl, etc.; R = H, alkyl; R1, R2 = H, halo, NO2, etc.; or R1 and R2 are linked together as OCH2O or OCH2CH2O] which are JAK3 kinase inhibitors, useful in treating asthma, host vs. graft rejection/transplantation or rheumatoid arthritis, were prep'd. E.g., a 7-step synthesis of I [X = NH; n = 0; Ar = 3-(hydroxymethyl)-2-methylphenyl; R1 = OCH2Ph; R2 = OMe], starting from 4-nitroguaiacol potassium salt, was given. The exemplified compds. I showed IC50 of < 25 .mu.M in JAK3 HTRF assay.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD



ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 8 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:594844 CAPLUS
 DN 137:140518
 TI Preparation of thiazolyl-, oxazolyl-, pyrrolyl-, and imidazolyl- acid amide derivatives as inhibitors of phosphodiesterase IV isozymes
 IN Marfat, Anthony; McKechney, Michael William
 PA Pfizer Products Inc., USA
 SO PCT Int. Appl., 249 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE	
PI	WO 2002060898	A1	20020808	WO 2001-IB2728	20011224	
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2001-265486PP	20010131
	US 2002123520	A1	20020905	US 2002-62145	20020131	
	US 6559168	B2	20030506	US 2001-265486PP	20010131	
	US 2003130254	A1	20030710	US 2002-300959	20021120	
				US 2001-265486PP	20010131	
				US 2002-62145	A320020131	
	US 2003186974	A1	20031002	US 2002-300950	20021120	
				US 2001-265486PP	20010131	
				US 2002-62145	A320020131	
OS	MARPAT 137:140518					
IT	68844-77-9 , Astemizole					
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)					
	(combination therapy with PDE4 inhibitors; prepn. of thiazolyl-, oxazolyl-, pyrrolyl-, and imidazolyl- acid amide derivs. as inhibitors of PDE4 isoenzymes)					
RN	68844-77-9 CAPLUS					
CN	1H-Benzimidazol-2-amine, 1-[(4-fluorophenyl)methyl]-N-[1-[2-(4-methoxyphenyl)ethyl]-4-piperidinyl]- (9CI) (CA INDEX NAME)					



GI



AB Title compds. I [wherein p = 0-1; q = 0-1; provided that when q = 0, n = 2; m = 0-3; n = 1-2; W1 and W2 = independently O, SOO-2, or NR3; or W2 = (un)substituted methylene; Y = SOO-2, O, NOO-1, NR3, or (un)substituted methylene; ; RA and RB = independently H, F, CF3, alkyl, or (un)substituted cycloalkyl, Ph, or benzyl; or when m = 1, CRARB = (un)substituted spiro; RC and RD have the same meaning as RA and RB except that one of them must be H; R1 and R2 = H, F, Cl, CN, NO2, (fluoro)alkyl, alkynyl, alkoxy, phenoxy, carbamoyl, etc.; R3 = H, alkyl, Ph, benzyl, alkoxy, phenoxy, etc.; R4, R5, and R6 = H, F, Cl, and (un)substituted (cyclo)alkyl, alkenyl, alkynyl, Ph, benzyl, pyridyl, alkoxy, phenoxy, acyl, carboxy, CN, NO2, carbamoyl, ureido, (hetero)aryl, etc.; G1 and G2 = independently (un)satd. carbocyclyl or heterocyclyl; E = (un)substituted carboxy, carbamoyl, acyl, hydroxyalkyl, cyanoalkyl, acylamino, ureido, amino, heterocyclyl, etc.] were prepd. as inhibitors of PDE4 (no data). For example, 4-(3-cyanophenoxy)thiazole-5-carboxylic acid was treated with 2-(4-aminomethylphenyl)propan-2-ol in the presence of EDCI and HOBT in DMF to give the thiazolamide II. I are useful in the treatment of diseases regulated by the activation and degranulation of eosinophils, esp. asthma, chronic bronchitis, and chronic obstructive pulmonary disease (no data). In addn., I may be used in combination therapy with a wide variety of other therapeutic agents.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 9 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2002:293652 CAPLUS

DN 136:325531
 TI Preparation of (poly)azanaphthalenyl carboxamides as HIV integrase inhibitors
 IN Anthony, Neville J.; Gomez, Robert P.; Young, Steven D.; Egbertson, Melissa; Wai, John S.; Zhuang, Linghang; Embrey, Mark; Tran, Lekhanh; Melamed, Jeffrey Y.; Langford, H. Marie; Guare, James P.; Fisher, Thorsten E.; Jolly, Samson M.; Kuo, Michelle S.; Perlow, Debra S.; Bennett, Jennifer J.; Funk, Timothy W.
 PA Merck & Co., Inc., USA
 SO PCT Int. Appl., 434 pp.
 CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002030930	A2	20020418	WO 2001-US31456	20011009
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2000-239707PP	20001012
				US 2001-281656PP	20010405
AU	2002011527	A5	20020422	AU 2002-11527	20011009
				US 2000-239707PP	20001012
				US 2001-281656PP	20010405
				WO 2001-US31456W	20011009
EP	1326865	A2	20030716	EP 2001-979582	20011009
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-239707PP	20001012
				US 2001-281656PP	20010405
				WO 2001-US31456W	20011009
US	2003055071	A1	20030320	US 2001-973853	20011010
				US 2000-239707PP	20001012
				US 2001-281656PP	20010405

PATENT FAMILY INFORMATION:



FAN 2002:293653

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002030931	A2	20020418	WO 2001-US42564	20011009
	WO 2002030931	A3	20021024		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2000-239707PP	20001012
				US 2001-281656PP	20010405

AU 2002011874	A5	20020422	AU 2002-11874	20011009
			US 2000-239707PP	20001012
			US 2001-281656PP	20010405
			WO 2001-US42564W	20011009
EE 200300145	A	20030616	EE 2003-145	20011009
			US 2000-239707PP	20001012
			US 2001-281656PP	20010405
			WO 2001-US42564W	20011009
US 2003055071	A1	20030320	US 2001-973853	20011010
			US 2000-239707PP	20001012
			US 2001-281656PP	20010405
NO 2003001672	A	20030605	NO 2003-1672	20030411
			US 2000-239707PP	20001012
			US 2001-281656PP	20010405
			WO 2001-US42564W	20011009
OS MARPAT 136:325531				
IT 410543-58-7P, 5-[(1-Benzylpiperidin-4-yl)amino]-N-(3,5-dichlorobenzyl)-8-hydroxy-1,6-naphthyridine-7-carboxamide				
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
			(HIV integrase inhibitor; prepn. of (poly)azanaphthalenyl carboxamides as HIV integrase inhibitors for treatment of AIDS)	
RN 410543-58-7 CAPLUS				
CN 1,6-Naphthyridine-7-carboxamide, N-[(3,5-dichlorophenyl)methyl]-8-hydroxy-5-[(1-(phenylmethyl)-4-piperidinyl)amino]- (9CI) (CA INDEX NAME)				



GI



AB Title compds., including certain quinoline carboxamide and naphthyridine carboxamide derivs., I [wherein A = (un)substituted Ph or Ph fused to a carbocycle; L = a single bond, or (un)substituted alkyl, alkenyl, alkylcycloalkylalkyl, or alkyl-M-alkyl; M = NR_a, OCO, or CO₂; X = N or CQ₁; Y = N or CQ₂, provided that X and Y are not both N; Z₁ = N or CQ₃; Z₂ = N or CQ₄; Z₃ = N or CH; Q₁-Q₄ = independently H, halo, CN, NR₁CR₁O, or (un)substituted alkyl, alkoxy, alkenyl, alkynyl, carbamoyl, carboximidamido, amino, etc.; or C₂Q₂Q₃ = (un)substituted 5- or 6-membered carbocycle or heterocycle; R₁ and R₂ = independently H, OH, halo, NO₂, CN, or (un)substituted alkyl, alkenyl, alkoxy, amino, sulfonylamino, etc.; R₃ and R₄ = independently H, halo, CN, NO₂, OH, alkenyl, or (un)substituted alkyl, amino, sulfonylamino, etc.; R₅ = H, CN, CN, or (un)substituted alkyl or aryl; Ra = independently H or (halo)alkyl; or pharmaceutically acceptable salts thereof] were prep'd. I are inhibitors of HIV integrase and inhibitors of HIV replication, and are useful in the prevention or treatment of infection by HIV and the treatment of AIDS, as compds. or pharmaceutically acceptable salts, or as ingredients in pharmaceutical compns., optionally in combination with other antivirals, immunomodulators, antibiotics, or vaccines. For example, Mitsunobu reaction of iso-Pr 3-(hydroxymethyl)pyridine-2-carboxylate with Me N-[(4-methylphenyl)sulfonyl]glycinate, followed by cyclization in the presence on NaOMe, afforded Me 8-hydroxy-1,6-naphthyridine-7-carboxylate. Coupling with 3,5-dichlorobenzylamine in toluene gave II. Representative compds. were assayed for the inhibition of acute HIV infection of T-lymphoid cells and demonstrated IC₉₅ values of < 20 .mu.M.

L10 ANSWER 10 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
AN 2002:240760 CAPLUS

DN 136:279470

TI Preparation of 6-[(substituted phenyl)methyl]quinoline and quinazoline derivatives as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases

IN Angibaud, Patrick Rene; Venet, Marc Gaston; Saha, Ashis Kumar; Mevellec, Laurence Anne

PA Janssen Pharmaceutica N.V., Belg.

SO PCT Int. Appl., 97 pp.


CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2002024683	A1	20020328	WO 2001-EP10895	20010918
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001093829	A5	20020402	EP 2000-203366 A	20000925
EP 1322636	A1	20030702	AU 2001-93829	20010918
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
			EP 2000-203366 A	20000925
OS MARPAT 136:279470			WO 2001-EP10895W	20010918
IT 406164-50-9P				
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (farnesyl transferase inhibitor; prepn. of quinoline and quinazoline derivs. as farnesyl transferase inhibitors for treatment of tumors and proliferative diseases)				
RN 406164-50-9 CAPLUS				
CN Tetrazolo[1,5-a]quinazoline-7-methanol, 5-(3-chlorophenyl)-.alpha.- (1- methyl-1H-imidazol-5-yl)-.alpha.- [4-[[[1-(phenylmethyl)-4- piperidinyl]amino]methyl]phenyl]- (9CI) (CA INDEX NAME)				



GI



AB Title compds. I [wherein m and n = independently 0-5; q = 0-3; Y1Y2 = C:N, C:CR9, CHNR9, or CHCHR9; C9 = H, halo, CN, (cyclo)alkyl, hydroxyalkyl, alkoxy(alkyl), aminoalkyl, (amino)alkenyl, (amino)alkynyl, halocarbonyl, hydroxycarbonyl, alkoxycarbonyl, aryl, (un)substituted amino or carbamoyl, etc.; R1 and R2 = independently azido, OH, halo, CN, NO₂, trihalomethyl, alkoxy, aryloxy, heterocyclyloxy, alkylthio, or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, carbamoyl, amino, sulfamoyl, etc.; or 2 adjacent R1 = OCH₂O, OCH₂CH₂O, OCH:CH, OCH₂CH₂, OCH₂CH₂CH₂, CH:CHCH:CH; R3 = H, halo, CN, alkenyl, alkynyl, hydroxycarbonyl, alkoxycarbonyl, aryl, heterocyclyl, alkoxy, alkylthio, (un)substituted (cyclo)alkyl or amino, etc.; R4 = (un)substituted imidazolyl, triazolyl, or pyridyl; R5 = CN, OH, halo, alkenyl, alkynyl, hydroxycarbonyl, alkoxycarbonyl, or (un)substituted (cyclo)alkyl, alkoxy, amino, or carbamoyl, etc.; R6 = halo or (un)substituted (cyclo)alkyl, alkenyl, alkynyl, alkylthio, carboxy, carbamoyl, acyl(amino), etc.; R7 = O or S; or R6R7 = (un)substituted CH:CHN:, CH>NN:, CONHN:, N>NN:, N:CHN:, CH:CHCH:, CH:NCH:, CONHCH:, N:NCH:, or CH₂(CH₂)₀₋₁CH₂N:; or pharmaceutically acceptable salts, N-oxides, or stereochem. isomeric forms thereof] were prep'd. For example, 6-bromo-2-chloro-4-(3-chlorophenyl)quinoline (6-step prepn. given) was coupled with 4-(diethoxymethyl)benzaldehyde in the presence of BuLi in THF to give the 6-quinolinemethanol (64%), which was treated with MnO₂ in 1,4-dioxane to afford the methanone. Methoxylation using MeONa in MeOH (74%), followed by addn. of 1-methyl-1H-imidazole in the presence of BULi and ClSiEt₃ in THF, gave 4-(3-chlorophenyl)-.alpha.-[4-(diethoxymethyl)phenyl]-2-methoxy-.alpha.-[(1-methyl-1H-imidazol-5-yl)-6-quinolinemethanol (56%). The latter was refluxed in HCl for 24 h, cooled, poured out into H₂O, and stirred at room temp. for 1 h to afford the quinolinone II.bul.HCl (98%). I have potent farnesyl transferase inhibitory effect and are useful for inhibiting proliferative diseases and growth of tumors expressing an activated ras oncogene (no data).

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 11 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:220574 CAPLUS

DN 136:263158

TI Benzimidazolyl-substituted quinolinone derivatives and analogs, with inhibitory action against vascular endothelial growth factor receptor tyrosine kinase, and useful as anticancer agents

IN Renhowe, Paul; Pecchi, Sabina; Machajewski, Tim; Shafer, Cynthia; Taylor, Clarke; McCrea, Bill; McBride, Chris; Jazan, Elisa; Wernette-Hammond, Mary-Ellen; Harris, Alex

PA Chiron Corporation, USA

SO PCT Int. Appl., 207 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022598	A1	20020321	WO 2001-US42131	20010911
	WO 2002022598	C1	20021121		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			US 2000-232159PP	20000911
	AU 2001093275	A5	20020326	AU 2001-93275	20010911
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
	EP 1317442	A1	20030611	EP 2001-973722	20010911
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-232159PP	20000911
				WO 2001-US42131W	20010911
	NO 2003001097	A	20030325	NO 2003-1097	20030310
				US 2000-232159PP	20000911
				WO 2001-US42131W	20010911

PATENT FAMILY INFORMATION:

FAN 2003:98039

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003028018	A1	20030206	US 2002-116117	20020405
				US 2000-232159PP	20000911
				US 2001-951265 A2	20010911
	US 2002107392	A1	20020808	US 2001-951265	20010911
	US 6605617	B2	20030812		
	US 2003158224	A1	20030821	US 2000-232159PP	20000911
				US 2002-284017	20021030
				US 2000-232159PP	20000911
				US 2001-951265 A1	20010911

OS MARPAT 136:263158


IT 405170-47-0P, 6-Chloro-3-(5-(morpholin-4-yl)-1H-benzimidazol-2-yl)-4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)-one
405170-62-9P, 6-Chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)piperidin-4-yl)amino]quinolin-2(1H)-one

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; prepn. of benzimidazolyl-substituted quinolinone derivs. and analogs as VEGFR tyrosine kinase-inhibiting anticancer agents)


RN 405170-47-0 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-morpholinyl)-1H-benzimidazol-2-yl]-4-[(1-(phenylmethyl)-4-piperidinyl)amino] - (9CI) (CA INDEX NAME)



RN 405170-62-9 CAPLUS

CN 2(1H)-Quinolinone, 6-chloro-3-[5-(4-methyl-1-piperazinyl)-1H-benzimidazol-2-yl]-4-[[1-(phenylmethyl)-4-piperidinyl]amino]- (9CI) (CA INDEX NAME)



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. of formulas I and II are provided [for I: Z = O, S, (un)substituted NH; Y = certain OH derivs., CHO, esters and amides of CO₂H, certain NH₂ derivs.; R₁-R₄ = H, halo, cyano, NO₂, OH or derivs., NH₂ or derivs., (un)substituted amidinyl, guanidinyl, alk(en/yn)yl, aryl, heterocyclyl, CHO, CO₂H and esters and amides; R₅-R₈ = H, halo, NO₂, OH or derivs., NH₂ or derivs., SH or derivs., cyano, etc.; R₉ = H, OH, (un)substituted alkoxy or aryloxy, NH₂ or derivs., (un)substituted alkyl or aryl, CHO, alkanoyl, aroyl; for II: A, B, D, E = C or N, with at least one being N; Y = H, OH or derivs., SH or derivs., NH₂ or derivs., cyano, various acyl groups, (un)substituted alk(en/yn)yl, aralkyl, heterocycloalkyl, aryl, etc.; R₁-R₈ = H, halo, NO₂, cyano, OH or derivs., NH₂ or derivs., acyl, SH or derivs., etc.; R₉ = H, OH, (un)substituted

alkoxy, aryloxy, NH₂ or derivs., aryl, CHO, alkanoyl, aroyl]. Also provided are pharmaceutical formulations including the compds. or their pharmaceutically acceptable salts and a pharmaceutically acceptable carrier, which may be prep'd. by mixing the compds. or salts with a carrier and water. A disclosed method of treating a patient includes administering a pharmaceutical formulation according to the invention to a patient. Claims include tautomers of the compds., pharmaceutically acceptable salts, and pharmaceutically acceptable salts of the tautomers. I and II are inhibitors of receptor tyrosine kinases, and particularly of vascular endothelial growth factor receptor (VEGFR) tyrosine kinase. As such, they are inhibitors of angiogenesis, and thereby act as anticancer agents. Approx 270 invention compds. are listed, with detailed preps. given for about 50 compds. Several general preparatory methods are discussed in detail. For instance, cyclocondensation of Et 2-(benzimidazol-2-yl)acetate with the corresponding ortho-amino nitrile (preps. given), carried out in refluxing ClCH₂CH₂Cl in the presence of SnCl₄, gave the invention quinolinone III. Many compds. I and II had in vitro IC₅₀ values of less than 10 .mu.M with respect to flt-1 (VEGFR1), KDR (VEGFR2) and bFGF kinases (recombinant, expressed in Sf9 insect cells).

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 12 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:185086 CAPLUS
 DN 136:247505
 TI Preparation of aminoquinolines as inhibitors of cGMP phosphodiesterase
 IN Bi, Yingzhi; Yu, Guixue; Rotella, David P.; Macor, John E.
 PA Bristol-Myers Squibb Company, USA
 SO PCT Int. Appl., 96 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002020489	A2	20020314	WO 2001-US26130	20010821
	WO 2002020489	A3	20020606		
				W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG	
				US 2000-230267PP	20000906
	US 2002177587	A1	20021128	US 2001-933066	20010820
	US 6576644	B2	20030610		
				US 2000-230267PP	20000906
	AU 2001085163	A5	20020322	AU 2001-85163	20010821
				US 2000-230267PP	20000906
				WO 2001-US26130W	20010821
OS	MARPAT 136:247505				
IT	403839-97-4P				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES				

(Uses)


(target compd.; prepn. of aminoquinolines as inhibitors of cGMP phosphodiesterase)

RN 403839-97-4 CAPLUS


CN 3-Quinolinecarboxamide, 4-[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



I



II

AB Title compds. I [R2, R6, R7, and R8 = independently H, halo,

(un)substituted alkyl, alkoxy, nitro, etc.; R4 and R5 = independently H, (un)substituted alkyl, cycloalkyl, aryl, or heteroaryl with provision R4 and R5 are not both H; R3 = (CH₂)_zY, wherein z = 0-3 and Y is independently selected from (un)substituted imidazole, triazole, OR₉, CO₂R₉, CH(CO₂R₉)₂, NR₁₀R₁₁, NR₁₀CONR₁₁R₁₂, etc.; or R4 and R5 together with Y form a heterocyclic ring; R9 = H, OH, (un)substituted alkyl, alkoxy, aryl, heteroaryl, etc.; R10, R11 and R12 = independently H, (un)substituted alkyl, alkoxy, cycloalkyl, heterocyclo, heteroaryl, etc.; or R10 forms a 3-7 membered heterocyclo ring with R11 or R12, or R11 forms a 3-7 membered ring with R12] are prepd. and disclosed as inhibitors of cGMP PDE, esp. type 5. Thus, II was prepd. via substitution of 4-chloro-6-cyanoquinoline-3-carboxylic acid Et ester with 3-chloro-4-methoxybenzylamine hydrochloride (97% yield). As inhibitors of cGMP phosphodiesterase, I are useful in treatment of cardiovascular disorders, diabetes, gastrointestinal disorders and sexual dysfunction, in particular erectile dysfunction (no data).

L10 ANSWER 13 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2002:123617 CAPLUS

DN 136:183819

TI Preparation of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors

IN Wang, Wei-Bo; Curtin, Michael L.; Fakhoury, Stephen A.; Gwaltney, Stephen L.; Hasvold, Lisa A.; Hutchins, Charles W.; Li, Qun; Lin, Nan-Horng; Nelson, Lissa Taka Jennings; O'Connor, Steve; Sham, Hing L.; Sullivan, Gerard M.; Wang, Gary T.; Wang, Xilu

PA USA

SO U.S. Pat. Appl. Publ., 189 pp.
 CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002019527	A1	20020214	US 2001-842391	20010425
				US 2000-200165PP	20000427

OS MARPAT 136:183819


IT 371761-79-4P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of (imidazolylalkyl)biphenylcarbonitriles and analogs as farnesyltransferase inhibitors)


RN 371761-79-4 CAPLUS

CN Benzonitrile, 4-[(1-methyl-1H-imidazol-5-yl)[[1-(phenylmethyl)-4-piperidinyl]amino)methyl]-2-(1-naphthalenyl)-, trihydrochloride (9CI) (CA INDEX NAME)



● 3 HCl

GI



AB Title compds. (I) were prep'd. Thus, 2-MeC₆H₄C₆H₃(CN)(CHO)-2,5 was condensed with 1-methyl-2-triethylsilyl-1H-imidazole (prepn. each given) and the product O-arylated to give title compd. II. Data for biol. activity of I were given.

L10 ANSWER 14 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:107318 CAPLUS

DN 136:151163

TI Preparation of indazole derivatives as JNK enzyme inhibitors

IN Bhagwat, Shripad S.; Satoh, Yoshitaka; Sakata, Steven T.

PA Signal Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 412 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002010137	A2	20020207	WO 2001-US23890	20010730
	WO 2002010137	C2	20030206		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO,

RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
US 2000-221799PP 20000731

US 2002103229 A1 20020801 US 2001-910950 20010723
US 2000-221799PP 20000731

EP 1313711 A2 20030528 EP 2001-957332 20010730
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
US 2000-221799PP 20000731
WO 2001-US23890W 20010730

OS MARPAT 136:151163

IT 395107-63-8P, N-[1-Benzyl-4-piperidyl]-3-[5-(1H-1,2,4-triazol-3-


yl)-1H-indazol-3-yl]benzamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
(Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
(Uses)

(prepn. of indazole derivs. as JNK enzyme inhibitors)

RN 395107-63-8 CAPLUS

CN Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]-3-[5-(1H-1,2,4-triazol-3-yl)-
1H-indazol-3-yl] - (9CI) (CA INDEX NAME)




AB Indazole derivs., 3-R1A-5-R2-1H-indazoles (1), having activity as selective inhibitors of JNK are disclosed. In 1: A is a direct bond, -(CH₂)_a, -(CH₂)_bCH:CH(CH₂)_c, or -(CH₂)_bC.tplbond.C(CH₂)_c; R1 is aryl, heteroaryl or heterocycle fused to Ph, each being optionally substituted with 1-4 R3; R2 is -R₃, -R₄, -(CH₂)_bC(O)R₅, -(CH₂)_bC(:O)OR₅, -(CH₂)_bC(O)NR₅R₆, -(CH₂)_bC(O)NR₅(CH₂)_cC(O)R₆, -(CH₂)_bNR₅C(O)R₆, -(CH₂)_bNR₅C(O)NR₆R₇, -(CH₂)_bNR₅R₆, -(CH₂)_bOR₅, -(CH₂)_bSODR₅ or -(CH₂)_bSO₂NR₅R₆. A is 1-6; b and c are the same or different and are 0-4; d is 0-2. R3 is at each occurrence independently halogen, hydroxy, carboxy, alkyl, alkoxy, haloalkyl, acyloxy, thioalkyl, sulfinylalkyl, sulfonylalkyl, hydroxyalkyl, aryl, substituted aryl, arylalkyl, substituted arylalkyl, heterocycle, substituted heterocycle, heterocyclealkyl, substituted heterocyclealkyl, -C(O)OR₈, -C(O)R₈, -C(O)NR₈R₉, -C(O)NR₈OR₉, -SO₂NR₈R₉, -NR₈SO₂R₉, -CN, -NO₂, -NR₈R₉, -NR₈C(O)R₉, -NR₈C(O)(CH₂)_bOR₉, -NR₈C(O)(CH₂)_bR₉, -O(CH₂)_bNR₅R₉, or heterocycle fused to Ph. R4 is alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, each being optionally substituted with 1-4 R3, or R4 is halogen or hydroxy. R5, R6 and R7 are the same or different and are H, alkyl, aryl, arylalkyl, heterocycle or heterocyclealkyl, wherein each of R5, R6 and R7 are optionally substituted with 1-4 R3. R8 and R9 are the same or different and at each occurrence independently H, alkyl, aryl, arylalkyl, heterocycle, or heterocyclealkyl, or R8 and R9 taken together with the atom or atoms to which they are bonded form a heterocycle, wherein each of R8, R9, and R8 and R9 taken together to form a heterocycle

are optionally substituted with 1-4 R3 with the proviso that: when A is a direct bond and R1 is Ph, R2 is not Me, methoxy, C(O)CH₃ or C(O)H; when A is a direct bond and R1 is 4-Me-Ph, R2 is not Me; when A is a direct bond and R1 is 4-F-Ph, R2 is not trifluoromethyl; when A is a direct bond or -C.tplbond.C- and R1 is Ph, R2 is not -COOEt; and when A is a direct bond and R1 is 6,7-dimethoxyisoquinolin-1-yl, R2 is not hydroxy. Such compds. have utility in the treatment of a wide range of conditions that are responsive to JNK inhibition. Thus, methods of treating such conditions are also disclosed, as are pharmaceutical compns. contg. one or more compds. of the above compds. Many of the claimed compds. have IC₅₀ values $\text{IC}_{50} \text{ req. } 0.5 \text{ } \mu\text{M}$ in the JNK2 assay, e.g. 5-[3-(4-fluorophenyl)-1H-indazol-5-yl]-2H-1,2,3,4-tetrazole. Although the methods of prepn. are not claimed, >400 example prepn.s. are included.

L10 ANSWER 15 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 2001:798200 CAPLUS
 DN 135:344482
 TI Preparation of substituted 4-(heteroaryl methyl)benzonitriles as farnesyltransferase inhibitors
 IN Wang, Wei-Bo; Curtin, Michael L.; Fakhoury, Stephen A.; Gwaltney, Stephen L., II; Hasvold, Lisa A.; Hutchins, Charles W.; Li, Qui; Lin, Nan-Horng; Jennings Nelson, Lissa Taka; O'Connor, Stephen J.; Sham, Hing L.; Sullivan, Gerald M.; Wang, Gary T.; Wang, Xilu
 PA Abbott Laboratories, USA
 SO PCT Int. Appl., 305 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001081316	A2	20011101	WO 2001-US13678	20010425
	WO 2001081316	A3	20020523		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			US 2000-563256 A 20000427
					US 2001-822205 A 20010402
EP	1276726	A2	20030122	EP 2001-932712	20010425
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			US 2000-563256 A 20000427
					US 2001-822205 A 20010402
					WO 2001-US13678W 20010425
OS	MARPAT 135:344482				
IT	371761-79-4P				
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(prepn. of substituted 4-(heteroaryl methyl)benzonitriles as farnesyltransferase inhibitors)				
RN	371761-79-4 CAPLUS				

CN Benzonitrile, 4-[(1-methyl-1H-imidazol-5-yl)[[1-(phenylmethyl)-4-piperidinyl]amino)methyl]-2-(1-naphthalenyl)-, trihydrochloride (9CI) (CA INDEX NAME)



● 3 HCl

GI



AB The title compds. [I; A1 = (un)substituted alkylene, etc.; R1 = halo, cycloalkyl, aryl, heteroaryl; R2 = heteroaryl selected from imidazolyl, pyrazolyl, pyrrolyl, etc.] and their pharmaceutically acceptable salts which farnesyltransferase, were prep'd. E.g., 3-step synthesis of the benzonitrile II.HCl which 88% inhibition of farnesyltransferase at 10⁻⁶ M, was given.

L10 ANSWER 16 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 2001:31473 CAPLUS

DN 134:100864

TI Indazole compounds and pharmaceutical compositions for inhibiting protein kinases, and methods for their use

IN Kania, Robert Steven; Bender, Steven Lee; Borchardt, Allen J.; Braganza,

John F.; Cripps, Stephan James; Hua, Ye; Johnson, Michael David; Johnson, Theodore Otto, Jr.; Luu, Hiep The; Palmer, Cynthia Louise; Reich, Siegfried Heinz; Tempczyk-russell, Anna Maria; Teng, Min; Thomas, Christine; Varney, Michael David; Wallace, Michael Brennan

PA Agouron Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 439 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001002369	A2	20010111	WO 2000-US18263	20000630
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				US 1999-142130PP 19990702
BR	2000012352	A	20020514	BR 2000-12352	20000630
				US 1999-142130PP	19990702
				WO 2000-US18263W	20000630
EP	1218348	A2	20020703	EP 2000-943375	20000630
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				US 1999-142130PP 19990702
				WO 2000-US18263W	20000630
JP	2003503481	T2	20030128	JP 2001-507809	20000630
				US 1999-142130PP	19990702
				WO 2000-US18263W	20000630
US	6531491	B1	20030311	US 2001-983786	20011025
				US 1999-142130PP	19990702
				US 2000-609335	B320000630
US	6534524	B1	20030318	US 2001-983783	20011025
				US 1999-142130PP	19990702
				US 2000-609335	B320000630
NO	2001005797	A	20020301	NO 2001-5797	20011128
				US 1999-142130PP	19990702
				WO 2000-US18263W	20000630
ZA	2001010061	A	20030206	ZA 2001-10061	20011206
				US 1999-142130PP	19990702
BG	106380	A	20020930	BG 2002-106380	20020201
				US 1999-142130PP	19990702
				WO 2000-US18263W	20000630
OS	MARPAT 134:100864				
IT	319466-31-4P				
	RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(prepn. of combinatorial libraries of aryl-substituted indazole derivs. as modulators and inhibitors of protein kinases in the treatment of tumor growth, cellular proliferation, and angiogenesis)				
RN	319466-31-4 CAPLUS				
CN	Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]-2-[(3-[(1E)-2-				

pyridinyl)ethenyl]-1H-indazol-6-yl]thio] - (9CI) (CA INDEX NAME)

Double bond geometry as shown.




GI




AB Indazole compds. I [R1 = substituted or unsubstituted aryl or heteroaryl, R₃CH:CH, R₃N:CH; R2 = substituted or unsubstituted aryl, heteroaryl, Y-X; R3 = substituted or unsubstituted alkyl alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl; Y = O, S, C(:CH₂), CO, SO, SO₂, alkylidene, NH, N(C₁-C₈ alkyl); X = substituted or unsubstituted aryl, heteroaryl, NH(alkyl), NH(cycloalkyl), NH(heterocycloalkyl), NH(aryl), NH(heteroaryl), NH(alkoxy), NH(dialkylamide)] and their pharmaceutically acceptable prodrugs, active metabolites, and salts are disclosed. The compds. modulate and/or inhibit the activity of certain protein kinases. In particular, I and pharmaceutical compns. contg. them are capable of mediating tyrosine kinase signal transduction, and thereby modulate and/or inhibit unwanted cell proliferation. The invention is also directed to the therapeutic or prophylactic use of pharmaceutical compns. contg. such compds., and to methods of treating cancer and other disease states assoc'd. with unwanted angiogenesis and/or cellular proliferation, such as diabetic retinopathy, neovascular glaucoma, rheumatoid arthritis, and psoriasis, by administering effective amts. of such compds. E.g., I [R1 = (E)-3,4-(MeO)2C₆H₃CH:CH; R2 = 4-HO-3-MeOC₆H₃] (II) was prep'd. from 6-aminoindazole by diazotization and substitution with iodide, protection of the indazole nitrogen with 2,4,6-Me₃C₆H₂SO₂Cl, coupling of the regioisomeric mixt. with 4-(methoxymethoxy)-3-methoxybenzeneboronic acid in the presence of dichlorobis(triphenylphosphine)palladium, and deprotection of the indazole moiety and iodination at the 3-position of the indazole. Treatment of the 3-indazolyl iodide with sec-butyllithium, phenyllithium, and DMF, regioselective protection of the indazole with 2,4,6-Me₃C₆H₂SO₂Cl, olefination with 3,4-dimethoxybenzyltriphenylphosphonium bromide, deprotection of the indazole, deprotection of the methoxymethyl group, and equilibration of the double bond with iodine gave II. Biol. data on protein kinase inhibition, cell proliferation

inhibition, neovascularization inhibition, and i.p. and oral bioavailability, are given.

L10 ANSWER 17 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1999:310758 CAPLUS
 DN 131:73602
 TI 4-N-linked-heterocyclic piperidine derivatives with high affinity and selectivity for human dopamine D4 receptors
 AU Moore, Kevin W.; Bonner, Katrine; Jones, Elizabeth A.; Emms, Frances; Leeson, Paul D.; Marwood, Rosemary; Patel, Shil; Patel, Smita; Rowley, Michael; Thomas, Steven; Carling, Robert W.
 CS Merck Sharp and Dohme Research Laboratories, Neuroscience Research Centre, Essex, CM20 2QR, UK
 SO Bioorganic & Medicinal Chemistry Letters (1999), 9(9), 1285-1290
 CODEN: BMCLE8; ISSN: 0960-894X
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 IT 971-34-6P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (prepn. of piperidinyl heterocycles and their affinity for dopamine D4 receptor)
 RN 971-34-6 CAPLUS
 CN Benzamide, N-[1-(phenylmethyl)-4-piperidinyl]- (9CI) (CA INDEX NAME)



GI



AB Several N-linked heterocyclic pyrazoles are prep'd. as hD4 ligands. The best compd. identified was I, which has high affinity for hD4 (5.2 nM) and >300-fold selectivity for hD4 receptors over hD2 and hD3 receptors.
 RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 18 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
 AN 1998:682229 CAPLUS

L10 ANSWER 18 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN
AN 1998:682229 CAPLUS

DN 129:302552
 TI Preparation of 1,4-disubstituted cyclic amine derivatives as serotonin antagonists
 IN Kitazawa, Noritaka; Ueno, Kohshi; Takahashi, Keiko; Kimura, Teiji; Sasaki, Atsushi; Kawano, Koki; Okabe, Tadashi; Komatsu, Makoto; Matsunaga, Manabu; Kubota, Atsuhiko
 PA Eisai Co., Ltd., Japan
 SO PCT Int. Appl., 635 pp.
 CODEN: PIXXD2
 DT Patent
 LA Japanese
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9843956	A1	19981008	WO 1998-JP1481	19980331
	W: AU, CA, CN, HU, JP, KR, MX, NO, NZ, RU, US			JP 1997-98433	A 19970331
	RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			JP 1997-366764	A 19971226
	AU 9865209	A1	19981022	AU 1998-65209	19980331
	AU 748038	B2	20020530	JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	ZA 9802707	A	19991020	WO 1998-JP1481	W 19980331
				ZA 1998-2707	19980331
	EP 976732	A1	20000202	JP 1997-98433	A 19970331
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE, FI			EP 1998-911137	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	NZ 337651	A	20020426	WO 1998-JP1481	W 19980331
				NZ 1998-337651	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	RU 2203275	C2	20030427	WO 1998-JP1481	W 19980331
				RU 1999-123039	19980331
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	US 6448243	B1	20020910	WO 1998-JP1481	W 19980331
				US 1999-367227	19990811
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	NO 9904720	A	19991130	WO 1998-JP1481	W 19980331
				NO 1999-4720	19990928
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
	US 2002086999	A1	20020704	WO 1998-JP1481	W 19980331
				US 2001-846259	20010502
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
				WO 1998-JP1481	W 19980331
	US 2002019531	A1	20020214	US 1999-367227	A319990811
	US 6579881	B2	20030617	US 2001-859517	20010518
				JP 1997-98433	A 19970331
				JP 1997-366764	A 19971226
				WO 1998-JP1481	W 19980331
				US 1999-367227	A319990811


OS MARPAT 129:302552

IT 202859-14-1P 214611-21-9P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (prepn. of 1,4-disubstituted cyclic amine derivs. as serotonin antagonists)


RN 202859-14-1 CAPLUS

CN 4-Piperidinamine, N-(3-methoxyphenyl)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 214611-21-9 CAPLUS

CN 4-Piperidinamine, N-(3-bromophenyl)-1-[2-(4-fluorophenyl)ethyl]- (9CI) (CA INDEX NAME)




IT 131587-28-5



RL: RCT (Reactant); RACT (Reactant or reagent)
 (prepn. of 1,4-disubstituted cyclic amine derivs. as serotonin antagonists)

RN 131587-28-5 CAPLUS

CN 4-Piperidinamine, N-(3-fluorophenyl)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



GI



AB The title compds. (I; A, B, C, D, T, Y, and Z each represents a methine group or a nitrogen atom; R1, R2, R3, R4, and R5 each represents a substituent, such as halo, OH, hydroxyalkoxy, lower alkyl, etc.; n is an integer of 0 to 3; m is an integer of 0 to 6; and p is an integer of 1 to 3; dotted bond represents a single or double bond) are prep'd. I have serotonin antagonism and serve as drugs for the treatment, alleviation and prevention of spastic paralysis or a central muscle relaxant for alleviating myotonia. Thus, indoline was reacted with 1-(4-fluorophenyl)-4-piperidone in the presence of NaB(OAc)₃ in AcOH and dichloroethane to give 63% the title compd. (II), which showed binding activity of 623.94 and > 200 nM for 5HT1a and 5HT2 resp.

RE.CNT 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10	ANSWER 19 OF 20	CAPLUS	COPYRIGHT 2003 ACS on STN
AN	1995:772553	CAPLUS	
DN	123:199300		
TI	Preparation of diaminopurinylribofuranuronamide derivatives as antiinflammatories.		
IN	Gregson, Michael; Ayres, Barry Edward; Ewan, George Blanch; Ellis, Frank; Knight, John		
PA	Glaxo Group Ltd., UK		
SO	PCT Int. Appl., 112 pp. CODEN: PIXXD2		
DT	Patent		
LA	English		
FAN.CNT 1			
	PATENT NO.	KIND	DATE
	-----	-----	-----
PI	WO 9417090	A1	19940804
	W: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN		WO 1994-EP145 19940118
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG		GB 1993-1000 A 19930120
CA	2153688	AA	19940804
			CA 1994-2153688 19940118
			GB 1993-1000 A 19930120
AU	9458851	A1	19940815
AU	679714	B2	19970710
			GB 1993-1000 A 19930120

ZA 9400335	A	19941024	WO 1994-EP145	W 19940118
			ZA 1994-335	19940118
			GB 1993-1000	A 19930120
EP 680488	A1	19951108	EP 1994-905100	19940118
EP 680488	B1	19980408		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
CN 1119440	A	19960327	CN 1994-191527	19940118
CN 1043997	B	19990707		
			GB 1993-1000	A 19930120
JP 08505864	T2	19960625	JP 1994-516652	19940118
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
AT 164849	E	19980415	AT 1994-905100	19940118
			GB 1993-1000	A 19930120
ES 2117249	T3	19980801	ES 1994-905100	19940118
			GB 1993-1000	A 19930120
RU 2129561	C1	19990427	RU 1995-122754	19940118
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
SK 281229	B6	20010118	SK 1995-918	19940118
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
IL 108372	A1	19980615	IL 1994-108372	19940119
			GB 1993-1000	A 19930120
FI 9503489	A	19950913	FI 1995-3489	19950719
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
NO 9502872	A	19950913	NO 1995-2872	19950719
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
US 5925624	A	19990720	US 1995-446727	19950918
			GB 1993-1000	A 19930120
			WO 1994-EP145	W 19940118
US 5889178	A	19990330	US 1997-934540	19970922
			GB 1993-1000	A 19930120
			US 1995-446727	A319950918

OS MARPAT 123:199300


IT 167297-77-0P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prep. of diaminopurinylribofuranuronamide derivs. as
 antiinflammatories)

RN 167297-77-0 CAPLUS

CN .beta.-D-Ribofuranuronamide, 1-deoxy-1-[6-[(2,2-diphenylethyl)amino]-2-[(1-phenylmethyl)-4-piperidinyl]amino]-9H-purin-9-yl]-N-ethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 167297-68-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (prep. of diaminopurinyleribofuranuronamide derivs. as antiinflammatories)


RN 167297-68-9 CAPLUS


CN .beta.-D-Ribofuranuronamide, 1-deoxy-1-[6-[(2,2-diphenylethyl)amino]-2-[(1-phenylmethyl)-4-piperidinyl]amino]-9H-purin-9-yl]-N-ethyl-2,3-O-(1-methylethylidene)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



GI





AB Title compds. [I; R₁ = H, C₃-8 cycloalkyl, C₁-6 alkyl; R₂ = (substituted) C₃-8 cycloalkyl, C₃-8 cycloalkyl-C₁-6 alkyl, pyrrolidin-3-yl, 2-oxopyrrolidin-4-yl, 2-oxopyrrolidin-5-yl, piperidin-3-yl, piperidin-4-yl, etc.; Q = O, S], were prep'd. Title compds. are useful as antiinflammatory agents, particularly in the treatment of patients with inflammatory conditions who are susceptible to leukocyte-induced tissue damage. Thus, (trans)-1-[2-[(4-aminocyclohexyl)amino]-6-[(2,2-diphenylethyl)amino]-9H-purin-9-yl]-1-deoxy-N-ethyl-2,3-O-(1-methylethylidene)-.beta.-D-ribofuranuronamide was stirred with aq. CF₃CO₂H to give (trans)-1-[2-[(4-aminocyclohexyl)amino]-6-[(2,2-diphenylethyl)amino]-9H-purin-9-yl]-1-deoxy-N-ethyl-.beta.-D-ribofuranonamide. The latter was 25 times more potent than NECA for inhibiting O₂- generation from neutrophils stimulated with fMLP, and inhibited ovalbumin-induced eosinophil accumulation in sensitized guinea pigs with ED₅₀ = 10 .mu.g/kg i.p.

L10 ANSWER 20 OF 20 CAPLUS COPYRIGHT 2003 ACS on STN

AN 1989:574103 CAPLUS

DN 111:174103

TI Preparation of piperidine-containing heterocycles as analgesics and anesthetics

IN Lin, Bor Sheng; Scheblein, Joseph W.

PA BOC Inc., USA

SO U.S., 28 pp.

CODEN: USXXAM

DT Patent

LA English


FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4831192	A	19890516	US 1987-139896	19871231
	EP 328830	A1	19890823	EP 1988-312149	19881221
	EP 328830	B1	19940601		
	R: BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE			US 1987-139896	19871231
				US 1987-139899	19871231
	ES 2054836	T3	19940816	ES 1988-312149	19881221
				US 1987-139896	19871231
				US 1987-139899	19871231
	JP 01301676	A2	19891205	JP 1988-332751	19881228
				US 1987-139896	19871231

PATENT FAMILY INFORMATION:

FAN 1989:423391


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4791120	A	19881213	US 1987-139899	19871231
	US 4871749	A	19891003	US 1988-256792	19881011
				US 1987-139899	19871231
	AU 8826604	A1	19890713	AU 1988-26604	19881206
	AU 616708	B2	19911107		
				US 1987-139899	19871231
	NO 8805463	A	19890703	NO 1988-5463	19881208
	NO 174553	B	19940214		
	NO 174553	C	19940525		
				US 1987-139899	19871231
IL 88645	A1	19930708	IL 1988-88645	19881209	
			US 1987-139899	19871231	
EP 328830	A1	19890823	EP 1988-312149	19881221	
EP 328830	B1	19940601			
R: BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE					
			US 1987-139896	19871231	
			US 1987-139899	19871231	
ES 2054836	T3	19940816	ES 1988-312149	19881221	
			US 1987-139896	19871231	
			US 1987-139899	19871231	
JP 01213278	A2	19890828	JP 1988-332752	19881228	
			US 1987-139899	19871231	
DK 8807328	A	19890701	DK 1988-7328	19881230	
			US 1987-139899	19871231	
FI 8806057	A	19890701	FI 1988-6057	19881230	
FI 92065	B	19940615			
FI 92065	C	19940926			
			US 1987-139899	19871231	
CN 1035285	A	19890906	CN 1989-100056	19881230	
CN 1023010	B	19931208			
			US 1987-139899	19871231	
OS CASREACT 111:174103; MARPAT 111:174103					
IT 968-86-5P 120070-52-2P 120070-54-4P					
120070-55-5P 120115-93-7P					
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)					
			(prepn. and reaction of, in prepn. of analgesics and anesthetics)		
RN 968-86-5 CAPLUS					
CN 4-Piperidinamine, 4-(phenylamino)-1-(phenylmethyl)- (9CI) (CA INDEX NAME)					



RN 120070-52-2 CAPLUS


CN 4-Piperidinamine, 4-(1-methyl-1H-1,2,4-triazol-5-yl)-N-phenyl-1-

(phenylmethyl) - (9CI) (CA INDEX NAME)




RN 120070-54-4 CAPLUS

CN 4-Piperidinamine, N-phenyl-1-(phenylmethyl)-4-(1H-tetrazol-5-yl)- (9CI) (CA INDEX NAME)




RN 120070-55-5 CAPLUS

CN 4-Piperidinamine, N-(2-fluorophenyl)-1-(phenylmethyl)-4-(1H-tetrazol-5-yl)- (9CI) (CA INDEX NAME)



RN 120115-93-7 CAPLUS


CN 4-Piperidinamine, 4-(4-methyl-2-thiazolyl)-N-phenyl-1-(phenylmethyl)- (9CI) (CA INDEX NAME)



IT 120070-56-6

RL: RCT (Reactant); RACT (Reactant or reagent)
(reaction of, in prepn. of analgesics and anesthetics)

RN 120070-56-6 CAPLUS
 CN 4-Piperidinecarbonitrile, 4-[(2-fluorophenyl)amino]-1-(phenylmethyl)-
 (9CI) (CA INDEX NAME)



GI For diagram(s), see printed CA Issue.
 AB Title compds. I [R1 = (substituted) oxadiazolyl, imidazolyl, triazolyl, tetrazolyl, thiazolyl; R2 = (substituted) Ph; R3 = acyl, alkoxycarbonyl; L = alkyl, alkoxy, thienylalkyl, (substituted) thiazolylalkyl, etc.] are prep'd. from I (R1 = cyano; R3 = H). Treatment of I (R1 = cyano; R2 = Ph; R3 = H; L = PhCH2) (prep'd. from KCN, PhNH2, and N-benzyl-4-piperidone, CAUTION: HCN evolution) with NaN3 in THF in the presence of AlCl3 gave I (R1 = 1H-tetrazol-5-yl), which was refluxed with Ac2O to afford I (R1 = 5-methyl-1,3,4-oxadiazol-2-yl; R2 = Ph; R3 = Ac; L = PhCH2). The oxalate of the latter showed ED50 >5.0 mg/kg in mice in a hot-plate analgesia test.

=> log y			
COST IN U.S. DOLLARS	SINCE FILE	TOTAL	
	ENTRY	SESSION	
FULL ESTIMATED COST	326.60	475.57	
 DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL	
	ENTRY	SESSION	
CA SUBSCRIBER PRICE	-35.81	-35.81	

STN INTERNATIONAL LOGOFF AT 11:01:01 ON 13 OCT 2003